



MMWRTM

Morbidity and Mortality Weekly Report

www.cdc.gov/mmwr

Surveillance Summaries

May 14, 2010 / Vol. 59 / No. SS-4

**Surveillance for Violent Deaths –
National Violent Death Reporting System,
16 States, 2007**

The *MMWR* series of publications is published by the Office of Surveillance, Epidemiology, and Laboratory Services, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

Suggested Citation: Centers for Disease Control and Prevention. [Title]. Surveillance Summaries, [Date]. MMWR 2010;59(No. SS-#).

Centers for Disease Control and Prevention

Thomas R. Frieden, MD, MPH
Director

Peter A. Briss, MD, MPH
Acting Associate Director for Science

James W. Stephens, PhD
Office of the Associate Director for Science

Stephen B. Thacker, MD, MSc
*Deputy Director for Surveillance,
Epidemiology, and Laboratory Services*

Editorial and Production Staff

Frederic E. Shaw, MD, JD
Editor, MMWR Series

Christine G. Casey, MD
Deputy Editor, MMWR Series

Teresa F. Rutledge
Managing Editor, MMWR Series

David C. Johnson
Lead Technical Writer-Editor

Douglas J. Bialecki
Project Editor

Martha F. Boyd
Lead Visual Information Specialist

Malbea A. LaPete
Stephen R. Spriggs

Terraye M. Starr
Visual Information Specialists

Quang M. Doan, MBA
Phyllis H. King

Information Technology Specialists

Editorial Board

William L. Roper, MD, MPH, Chapel Hill, NC, Chairman

Virginia A. Caine, MD, Indianapolis, IN

Jonathan E. Fielding, MD, MPH, MBA, Los Angeles, CA

David W. Fleming, MD, Seattle, WA

William E. Halperin, MD, DrPH, MPH, Newark, NJ

King K. Holmes, MD, PhD, Seattle, WA

Deborah Holtzman, PhD, Atlanta, GA

John K. Iglehart, Bethesda, MD

Dennis G. Maki, MD, Madison, WI

Patricia Quinlisk, MD, MPH, Des Moines, IA

Patrick L. Remington, MD, MPH, Madison, WI

Barbara K. Rimer, DrPH, Chapel Hill, NC

John V. Rullan, MD, MPH, San Juan, PR

William Schaffner, MD, Nashville, TN

Anne Schuchat, MD, Atlanta, GA

Dixie E. Snider, MD, MPH, Atlanta, GA

John W. Ward, MD, Atlanta, GA

CONTENTS

Introduction	2
Methods	3
Results	8
Discussion.....	13
Conclusion.....	16
References	16

Surveillance for Violent Deaths – National Violent Death Reporting System, 16 States, 2007

Debra L. Karch, PhD
Linda L. Dahlberg, PhD
Nimesh Patel, MS

Division of Violence Prevention, National Center for Injury Prevention and Control, CDC

Abstract

Problem/Condition: An estimated 50,000 persons die annually in the United States as a result of violence-related injuries. This report summarizes data from CDC's National Violent Death Reporting System (NVDRS) regarding violent deaths from 16 states for 2007. Results are reported by sex, age group, race/ethnicity, marital status, location of injury, method of injury, circumstances of injury, and other selected characteristics.

Reporting Period Covered: 2007.

Description of System: NVDRS collects data regarding violent deaths obtained from death certificates, coroner/medical examiner reports, and law enforcement reports. NVDRS began operation in 2003 with seven states (Alaska, Maryland, Massachusetts, New Jersey, Oregon, South Carolina, and Virginia) participating; six states (Colorado, Georgia, North Carolina, Oklahoma, Rhode Island, and Wisconsin) joined in 2004, four (California, Kentucky, New Mexico, and Utah) in 2005, and two states (Ohio and Michigan) were funded to begin data collection in 2010, totaling 19 states. This report includes data from 16 states that collected statewide data in 2007. California data are not included in this report because NVDRS data are collected only in a limited number of California cities and counties rather than statewide. Ohio and Michigan are excluded because they did not begin data collection until 2010.

Results: For 2007, a total of 15,882 fatal incidents involving 16,319 deaths occurred in the 16 NVDRS states included in this report. The majority (56.6%) of deaths was suicides, followed by homicides and deaths involving legal intervention (i.e., deaths caused by police and other persons with legal authority to use deadly force, excluding legal executions) (28.0%), deaths of undetermined intent (14.7%), and unintentional firearm deaths (0.7%). Suicides occurred at higher rates among males, American Indians/Alaska Natives, non-Hispanic whites, and persons aged 45–54 years. Suicides occurred most often in a house or apartment and involved the use of firearms. Suicides were precipitated primarily by mental-health, intimate-partner, or physical-health problems, or by a crisis during the preceding 2 weeks. Homicides occurred at higher rates among males and persons aged 20–24 years; rates were highest among non-Hispanic black males. The majority of homicides involved the use of a firearm and occurred in a house or apartment or on a street/highway. Homicides were precipitated primarily by arguments and interpersonal conflicts or in conjunction with another crime. Other manners of death and special situations or populations also are highlighted in this report.

Interpretation: This report provides a detailed summary of data from NVDRS for 2007. The results indicate that violent deaths resulting from self-inflicted or interpersonal violence disproportionately affected adults aged <55 years, males, and certain minority populations. For homicides and suicides, relationship problems, interpersonal conflicts, mental-health problems, and recent crises were among the primary precipitating factors. Because additional information might be reported subsequently as participating states update their findings, the data provided in this report are preliminary.

Public Health Action: For the occurrence of violent deaths in the United States to be better understood and ultimately prevented, accurate, timely, and comprehensive surveillance data are necessary. NVDRS data can be used to monitor the occurrence of violence-related fatal injuries and assist public health authorities in the development, implementation, and evaluation of programs and policies to reduce and prevent violent deaths at the national, state, and local levels. The continued development and expansion of NVDRS is essential to CDC's efforts to reduce the personal, familial, and societal costs of violence. Further efforts are needed to increase the number of states participating in NVDRS, with an ultimate goal of full national representation.

Corresponding Author: Debra L. Karch, PhD, Division of Violence Prevention, National Center for Injury Prevention and Control, CDC, 4770 Buford Hwy, N.E., MS F-64, Atlanta, GA, 30341-3724. Telephone: 770-488-1307; Fax: 770-488-4222; E-mail: dkarch@cdc.gov.

Introduction

An estimated 50,000 persons die annually in the United States from violence-related injuries. Homicide is the second leading cause of death for persons aged 15–24 years, the third leading cause for persons aged 10–14 and 25–34 years, and the fourth leading cause for persons aged 1–9 years. Suicide is the second leading cause of death for persons aged 25–34 years, the third leading cause for persons aged 15–24 years, and the fourth leading cause for persons aged 10–14 and 35–44 years. Only unintentional injury in those aged 1–34 years and malignant neoplasms and congenital anomalies in children aged 1–14 years were more common (1).

Public health authorities require accurate, timely, and comprehensive surveillance data to better understand and ultimately prevent the occurrence of violent deaths in the United States (2). In 2000, CDC started planning to implement the National Violent Death Reporting System (NVDRS) (3,4). The goals of this system are to:

- collect and analyze timely, high-quality data that monitor the magnitude and characteristics of violent death at the national, state, and local levels;
- ensure that data are disseminated routinely and expeditiously to public health officials, law enforcement officials, policy makers, and the public;
- ensure that data are used to develop, implement, and evaluate programs and policies that are intended to reduce and prevent violent deaths and injuries at the national, state, and local levels; and
- build and strengthen partnerships among organizations and communities at the national, state, and local levels to ensure that data are collected and used to reduce and prevent violent deaths and injuries.

NVDRS was conceived as a state-based active surveillance system that would collect risk-factor data concerning all violence-related deaths, including homicides, suicides, and legal intervention deaths (i.e., deaths caused by police and other persons with legal authority to use deadly force, excluding legal executions) as well as unintentional firearm deaths and deaths of undetermined intent. NVDRS data are used to assist the development, implementation, and evaluation of programs and policies designed to reduce and prevent these deaths and injuries at the national, state, and local levels.


Before implementation of NVDRS, single data sources (e.g., death certificates or supplemental homicide reports) provided limited information and circumstances from which to understand patterns of deaths collected by this system. NVDRS fills

this gap in national surveillance; it is the first system to provide detailed information on circumstances precipitating violent deaths, the first to link multiple source documents to enable researchers to understand each death more completely, and the first to link multiple deaths that are related to one another (e.g., multiple homicides, suicide pacts, and cases of homicide followed by the suicide of the suspected perpetrator).

NVDRS began operation in 2003 with seven states (Alaska, Maryland, Massachusetts, New Jersey, Oregon, South Carolina, and Virginia) participating; six states (Colorado, Georgia, North Carolina, Oklahoma, Rhode Island, and Wisconsin) joined in 2004, four (California, Kentucky, New Mexico, and Utah) in 2005, and two states (Ohio and Michigan) were funded to begin data collection in 2010, totaling 19 states (Figure). CDC provides funding for state participation and anticipates that NVDRS will expand to include all 50 states, the District of Columbia, and U.S. territories.

This report summarizes data for 2007 concerning deaths meeting NVDRS inclusion criteria from 16 states that collected statewide data (approximately 26% of the U.S. population). California data are not included in this report because NVDRS data are collected only in a limited number of California cities and counties rather than statewide. Ohio and Michigan were excluded because they did not begin data collection until 2010. Because additional information might be reported subsequently as participating states update their findings, the data provided in this report are preliminary. Annual updates of NVDRS data also are available through a web-based query system at <http://wisqars.cdc.gov:8080/nvdrs/nvdrsDisplay.jsp>.

FIGURE. States participating in the National Violent Death Reporting System, by year of initial data collection — United States, 2003–2010



Methods

NVDRS uses multiple, complementary data sources including death certificates, coroner/medical examiner (CME) records, and law enforcement reports. Some participating states use secondary sources (e.g., child fatality review team data, supplementary homicide reports, hospital data, and crime laboratory data). Bureau of Alcohol, Tobacco, Firearms, and Explosives traces information concerning firearms. NVDRS links together multiple documents for each violent death and also links multiple deaths that are related to each other (e.g., multiple homicides, a homicide followed by a suicide, or multiple suicides) into a single incident. The ability to analyze data linked in this way permits a comprehensive assessment of violent deaths.

NVDRS defines a violent death as a death resulting either from the intentional use of physical force or power against oneself, another person, or a group or community. In addition, NVDRS collects information regarding unintentional firearm injury deaths (i.e., incidents in which the person causing the injury did not intend to discharge the firearm) and deaths of undetermined intent. NVDRS case definitions are coded on the basis of the *International Classification of Diseases, 10th Revision* (ICD-10) (5). Cases with selected ICD-10 codes are included in NVDRS (Box 1). ICD-10 case finding is completed by participating states.

Variables analyzed in NVDRS include the following:

- manner of death (i.e., the intent of the person inflicting a fatal injury);
- mechanism of injury (i.e., the method used to inflict a fatal injury);
- circumstances preceding injury (i.e., the precipitating events that led to the infliction of a fatal injury);
- whether the decedent was a victim (i.e., a person who died because of a violence-related injury);
- whether the decedent was a suspect (i.e., a person believed to have inflicted a fatal injury on a victim);

- whether the decedent was both a suspect and a victim (i.e., a person believed to have inflicted a fatal injury on a victim and then was fatally injured himself or herself);
- incident (i.e., an occurrence in which one or more persons sustained a fatal injury that was linked to a common event during a 24-hour period); and
- type of incident (i.e., a combination of the manner of death and the number of victims in an incident).

NVDRS is incident-based, and all decedents (both victims and alleged perpetrators [suspects]) associated with a given incident are grouped in one record. Decisions about whether two or more deaths belong to the same incident are made on the basis of the timing of the injuries rather than on that of the deaths. Examples of a violent death incident include 1) a single isolated violent death, 2) two or more related homicides (including legal interventions) when the fatal injuries were inflicted <24 hours apart, 3) two or more related suicides or deaths of undetermined intent when the fatal injuries were inflicted <24 hours apart, and 4) a homicide followed by a related suicide when both fatal injuries were inflicted <24 hours apart.

Data are obtained from individual information sources and entered into source-specific computerized data entry screens (i.e., police report data are entered into police report screens and death certificate data into death certificate screens). In addition to allowing independent entry of each source, this approach permits later review of what each source contributed and identification of missing sources. This permits comparisons of the quality and completeness of state-specific data sources and allows states to provide feedback to sources regarding the consistency of their data compared with data from other sources. In addition, the system permits automatic electronic importation of specific data sources without requiring manual entry.

Abstraction of identical variables across multiple source documents can result in data inconsistencies, which NVDRS resolves by assigning a primacy (i.e., hierarchical) rule for each

BOX 1. International Classification of Diseases, Tenth Revision (ICD-10) codes used in the National Violent Death Reporting System

Manner of death	Death ≤1 year after injury	Death >1 year after injury
Intentional self-harm (suicide)	X60–X84	Y87.0
Assault (homicide)	X85–X99, Y00–Y09	Y87.1
Event of undetermined intent	Y10–Y34	Y87.2, Y89.9
Unintentional exposure to inanimate mechanical forces (firearms)	W32–W34	Y86 determined to be attributable to firearms
Legal intervention excluding executions, (Y35.5)	Y35.0–Y35.4, Y35.6–Y35.7	Y89.0
Terrorism	U01, U03	U02

variable. The primacy rules are applied to create a final analysis data set that uses data from all available sources. For each variable in NVDRS, primacy is established on the basis of a hierarchy of assumed reliability of all the sources for a single variable. For example, sex is collected in all three required documents (death certificate, CME record, and police report). The primacy for sex is expressed as death certificate/CME record/police report, which means the analysis file is constructed using the sex recorded in the death certificate; if this is left blank or is unknown, the sex recorded in the CME record is used; and if the CME record does not provide the sex or lists the sex as unknown, the police report is used.

Manner of Death

A manner (i.e., intent) of death for each decedent is assigned by a trained abstractor who takes into account information from all source documents. Typically, these documents are consistent regarding the manner of death, and the abstractor-assigned manner of death corresponds to that reported in all the source documents. On rare occasions, when a discrepancy exists among the source documents, the abstractor must assign a manner of death on the basis of the preponderance of evidence in the source documents. For example, if two sources classify a death as a suicide and a third classifies it as undetermined, the death will be coded as a suicide.

NVDRS classifies data using one of five abstractor-assigned manners of death:

- **Suicide.** Suicide is defined as a death resulting from the use of force against oneself when a preponderance of the evidence indicates that the use of force was intentional. This category includes deaths of persons who intended only to injure rather than kill themselves, deaths associated with risk taking behavior without clear intent to inflict fatal injury but associated with high risk of death (e.g., Russian roulette) and suicides involving only passive assistance to the decedent (e.g., supplying the means or information needed to complete the act). The category does not include deaths caused by chronic or acute substance abuse without the intent to die or deaths attributed to autoerotic behavior (e.g., self-strangulation during sexual activity). Corresponding ICD-10 codes included in NVDRS are X60–X84 and Y87.0.
- **Homicide.** Homicide is defined as a death resulting from the use of physical force or power, threatened or actual, against another person, group, or community when a preponderance of evidence indicates that the use of force was intentional. Two special scenarios that the National Center for Health Statistics (NCHS) regards as homicides are included in the NVDRS definition: 1) arson with no

intent to injure a person and 2) a stabbing with intent unspecified. This category excludes vehicular homicide without intent to injure, unintentional firearm deaths (a separate category listed below), combat deaths or acts of war, and deaths of unborn fetuses. Corresponding ICD-10 codes included in NVDRS are X85–X99, Y00–Y09, and Y87.1.

- **Unintentional firearm.** The term “unintentional firearm death” is used when a death results from a penetrating injury or gunshot wound from a weapon that uses a powder charge to fire a projectile and for which a preponderance of evidence indicates that the shooting was not directed intentionally at the decedent. Examples of deaths included in this category include the death of a person as a result of celebratory firing that was not intended to frighten, control, or harm anyone; a soldier shot during a field exercise but not in a combat situation; and a person who received a self-inflicted wound while playing with a firearm. This category excludes firearm injuries caused by unintentionally striking a person with the firearm (e.g., hitting a person on the head with the firearm rather than firing a projectile) and unintentional injuries from nonpowder guns (e.g., BB, pellet, or other compressed air- or gas-powered guns). Corresponding ICD-10 codes included in NVDRS are W32–W34 and Y86 with a method of firearm.
- **Undetermined intent.** The term “undetermined intent” is used when a death results from the use of force or power against oneself or another person for which the evidence indicating one manner of death is no more compelling than evidence indicating another. This category includes CME rulings (e.g., accident or suicide, undetermined, jumped or fell, self-inflicted injuries) when records give no evidence or opinions in favor of either unintentional or intentional injury. Corresponding ICD-10 codes included in NVDRS are Y10–Y34, Y87.2, and Y89.9.
- **Legal intervention.** The term “legal intervention” is used when a decedent is killed by a police officer or other peace officer (a person with specified legal authority to use deadly force), including military police, acting in the line of duty. This category excludes legal executions. Corresponding ICD-10 codes included in NVDRS are Y35.0–Y35.4, Y35.6, Y35.7, and Y89.0.

Comparability of NVDRS Surveillance Summary Data Prior to 2006

Four changes were made to how variables were reported between 2005 and 2006 that affect their comparability; no variable changes were made between 2006 and 2007. The 2005

to 2006 changes involve race/ethnicity, location of injury, relationship of victim to suspect, and method of injury. In 2005, the race variable was reported in six categories (white, black, Asian Pacific Islander [API], American Indian/Alaska Native [AI/AN], other, and unknown). Ethnicity was categorized separately as persons of any race that reported Hispanic origin. When this methodology was used, Hispanics were reported both within their race category and then again separately by ethnicity. The 2006 and 2007 methodology classifies each person as non-Hispanic white, non-Hispanic black, API, AI/AN, Hispanic, other, and unknown. Race and ethnicity are combined in one variable. This change allows for better comparability with other violence-related data.

Location of injury is coded from a list of 31 location options in NVDRS. Because certain options are selected rarely, certain response categories have been combined. In 2006 and 2007, the category “bank” was included in “office building” rather than in “commercial/retail area” as it was in 2005. Also in 2006 and 2007, the category “synagogue/church/temple” was subsumed under “other” and not reported separately as in 2005.

Relationship of the victim to the suspect includes a new category, “other intimate-partner involvement,” to refer to a death that is intimate-partner-related but that does not occur between the intimate partners themselves (e.g., when a child is killed by a parent’s partner). In addition, the categories “rival gang member” and “victim was injured by a law enforcement officer” are reported in 2006 and 2007 as separate categories; in 2005, these categories were included in “other specified relationship.” The categories “foster child” and “foster parent” also were moved from “other relative” to “child” and “parent,” respectively.

Four new categories were added to method of injury in 2006 and 2007: “firearm and poisoning,” “firearm and other method type,” “poisoning and other method type,” and “other combination of methods.” All deaths in these new categories involved more than one method, and the evidence did not indicate which method caused the fatal injury. For example, a homicide victim might have injuries from both a firearm and a sharp instrument, but the method that actually caused the fatal injury might be unclear. In this case, the method of injury would be categorized as “firearm and other method.”

Variables Analyzed

NVDRS collects approximately 250 unique variables (available at <http://www.cdc.gov/violenceprevention/nvdrs/index.htm>). The number of variables recorded for each incident depends on the content and completeness of the source documents. Variables include manner of death, demographics, ICD-10 and underlying cause-of-death codes and text, loca-

tion and date/time of injury and death, toxicology results, bodily injuries, precipitating circumstances, decedent-suspect relationship, and method of injury (Boxes 2 and 3).

Circumstances Preceding Death

The circumstances preceding death are defined as the precipitating events that led to the infliction of a fatal injury (Box 3). The circumstances that preceded a fatal injury are reported on the basis of the content of the CME record and police reports. Different sets of circumstances are coded for suicide/undetermined deaths, homicide/legal-intervention deaths, and unintentional firearm deaths. The variable “circumstances known” is a gateway variable to a list of potential circumstances. Each incident requires the data abstractor to code all circumstances in cases for which the circumstances are known. If circumstances are not known (e.g., for a body found in the woods with no other detail) the data abstractor leaves the gateway variable blank, and these cases are excluded from the denominator for circumstance values. If either the CME record or the police report indicates that the circumstance is reported to be true, then the abstractor enters data as confirmed (e.g., if the police report indicated that a decedent had disclosed an intent to commit suicide, then suicidal intent is accepted to be true).

BOX 2. Methods of injury — National Violent Death Reporting System, 16 states, 2007

- Firearm: method that uses a powder charge to fire a projectile.
- Sharp instrument: knife, razor, machete, pointed instrument (e.g., chisel or broken glass).
- Blunt instrument: club, bat, rock, or brick.
- Poisoning: street drug, alcohol, pharmaceutical, carbon monoxide, gas, rat poison, or insecticide.
- Hanging/strangulation/suffocation: hanging by the neck, manual strangulation, or plastic bag over the head.
- Personal weapons: hands, fists, or feet.
- Fall: being pushed or jumping.
- Drowning: inhalation of liquid in bathtub, lake, or other source of water/liquid.
- Fire/burn: inhalation of smoke or the direct effects of fire or chemical burns.
- Shaking: shaking a baby, child, or adult.
- Motor vehicle: car, bus, or motorcycle.
- Other transport vehicle: train or airplane.
- Intentional neglect: starvation, lack of adequate supervision, or withholding of health care.
- Other: any method other than those listed above.
- Unknown: method not reported or not known.

BOX 3. Circumstances preceding fatal injury, by manner of death — National Violent Death Reporting System, 16 states, 2007**Suicide/Undetermined Intent**

- Current depressed mood: decedent was perceived by self or others to be depressed.
- Current mental health problem: decedent has been identified as having a mental health disorder or syndrome listed in the Diagnostic and Statistical Manual, Version IV (DSM-IV).
- First/second type of mental illness diagnosis: identifies the DSM-IV diagnosis made by a medical or mental health practitioner.
- Current treatment for mental illness: decedent was currently receiving mental health treatment as evidenced by a current psychotropic medication or visit to a mental health professional in the previous 2 months.
- Alcohol/other substance problem: decedent was perceived by self or others to have a problem with, or to be addicted to, alcohol or other drugs.
- Person left a suicide note: decedent left a note, e-mail message, video, or other communication indicating an intent to die by suicide.
- Disclosed intent to die by suicide: decedent had previously expressed suicidal feelings to another person with time for that person to intervene; disclosure only at the time of the event, with no opportunity to intervene, is not coded as “disclosed intent to commit suicide.”
- History of suicide attempts: decedent was known to have made previous attempts, regardless of the severity of those attempts.
- Crisis during previous 2 weeks: a very current crisis or acute precipitating event appears to have contributed to the suicide. This is designed to measure impulsivity. The crisis event must have occurred in the previous 2 weeks or be impending in the following 2 weeks (e.g., a trial for a criminal offense begins the following week).
- Physical health problem: decedent was experiencing physical health problems that are believed to have contributed to the suicide (e.g., a recent cancer diagnosis or chronic pain).
- Intimate partner problem: problems with a current or former intimate partner that appear to have contributed to the suicide.
- Other relationship problem: problems with a family member, friend, or associate (other than an intimate partner) that appear to have contributed to the suicide.
- Job problem: decedent was either experiencing a problem at work or was having a problem with joblessness.
- School problem: decedent was experiencing a problem such as poor grades, bullying, social exclusion at school, or performance pressures.
- Financial problem: decedent was experiencing problems such as bankruptcy, overwhelming debt, or foreclosure of a home or business.
- Suicide of friend or family in previous 5 years: decedent was distraught over, or reacting to, a relatively recent suicide of a friend or family member.
- Other death of friend or family in previous 5 years: decedent was distraught over, or reacting to, a relatively recent nonsuicide death of a friend or family member.
- Recent criminal legal problem: decedent was facing criminal legal problems that appear to be associated with the suicide.
- Other legal problem: decedent was facing civil legal problems (e.g., a child custody or civil lawsuit).
- Perpetrator of interpersonal violence in previous month: decedent perpetrated interpersonal violence (e.g., being sought by police for assault or having been issued a restraining order resulting from recent violence) during the previous month.
- Victim of interpersonal violence in previous month: decedent was the target of interpersonal violence in the past month.

Homicide/Legal Intervention

- Precipitated by another crime: incident occurred as the result of another serious crime.
- Nature of crime: identifies the actual crime (e.g., robbery or drug trafficking).
- Crime in progress: crime was in progress at the time of the death.
- Argument over money/property: conflict between decedent and suspect was over money or property (including drugs).
- Other argument, abuse, conflict: conflict between decedent and suspect was over something other than money, property, or drugs.
- Jealousy (“lovers’ triangle”): jealousy or distress over an intimate partner’s relationship or suspected relationship with another person led to the homicide.
- Intimate-partner violence–related: homicide is related to conflict between current or former intimate partners; includes the death of actual intimate partners and nonintimate partner decedents killed to cause pain to an intimate partner (e.g., child or parent).

BOX 3. (Continued) Circumstances preceding fatal injury, by manner of death — National Violent Death Reporting System, 16 states, 2007

- Drug involvement: drug dealing or illegal drug use is suspected to have played a role in precipitating the homicide.
- Gang-related: homicide is suspected to have resulted from gang activity or gang rivalry; not used if the decedent was a gang member but the homicide did not appear to result from gang activity.
- Hate crime: decedent was intentionally selected because of his/her actual or perceived gender, religion, sexual orientation, race/ethnicity, or disability.
- Brawl: mutual physical fight involving three or more persons.
- Decedent was a bystander: decedent was not directly involved in the incident.
- Decedent was a police officer on duty: a law enforcement officer killed in the line of duty.
- Decedent was an intervener assisting a crime victim: decedent was attempting to assist a crime victim at the time of the incident (e.g., a child attempts to intervene and is killed while trying to assist a parent who is being assaulted).
- Mercy killing: the decedent wished to die because of terminal or hopeless disease or condition, and documentation indicates that the decedent wanted to be killed.
- Celebratory firing: shooter fired the gun upward in a celebratory manner with no intention of threatening or endangering others.
- Loading/unloading gun: firearm discharged when the shooter was loading/unloading ammunition.
- Cleaning gun: firearm discharged when the shooter was cleaning the gun.
- Showing gun to others: showing the gun to another person when the gun discharged or the trigger was pulled.
- Playing with gun: the shooter and one or more others were playing with a gun.
- Thought safety was engaged: shooter thought the gun was inoperable because the safety was engaged.
- Thought unloaded/magazine disengaged: shooter thought the gun was unloaded because the magazine was disengaged.
- Thought gun was unloaded/other: shooter thought the gun was unloaded for other unspecified reason.
- Unintentionally pulled trigger: shooter unintentionally pulled the trigger (e.g., while grabbing the gun or holding it too tightly).
- Bullet ricochet: bullet ricocheted from its intended target and unintentionally struck the decedent.
- Gun defect or malfunction: gun had a defect or malfunctioned as determined by a trained firearm examiner.
- Fired while holstering/unholstering: gun was being replaced or removed from holster/clothing.
- Dropped gun: gun discharged when it was dropped or when something was dropped on it.
- Fired while operating safety/lock: shooter unintentionally fired the gun while operating the safety lock.
- Gun mistaken for toy: gun was mistaken for a toy and was fired without the user understanding the danger.

Unintentional Firearm Death

- Hunting: death occurred anytime after leaving home for a hunting trip and before returning home from a hunting trip; the shooting need not have been during an active hunt to be coded.
- Target shooting: a shooter was aiming for a target and unintentionally hit a person; can be at a shooting range or an informal backyard setting.
- Self-defensive shooting: self-inflicted shooting in which the decedent was attempting to use a gun in self-defense.

Coding Training and Quality Control

Coding training is held annually for all participating states. Ongoing coding support is provided through an e-mail help desk, monthly conference calls with all states, and regular conference calls with individual states. A coding manual is provided. Software features enhance coding reliability, including automated validation rules and a hover-over feature containing variable-specific information. Details regarding NVDRS procedures and coding are available at <http://www.cdc.gov/violenceprevention/nvdrs/publications.htm>.

States are requested to perform blind reabstraction of cases using multiple abstractors to identify inconsistencies. CDC also runs a quality-control analysis in which multiple variables are reviewed for their appropriateness, with special focus on abstractor-assigned variables (e.g., method selection and manner of death). If CDC questions any variable, CDC notifies the state for a response or correction.

Time Frame

States are required to report all deaths within 6 months of the end of each calendar year for the preceding January–December time frame. States then have an additional 12 months to complete each incident record. Although states typically meet these timelines, additional details sometimes arrive after a deadline has passed. New incidents also might be identified after the deadline (e.g., if a death certificate is revised, new evidence is obtained that changes a manner of death, or a miscoded ICD-10 is corrected to meet NVDRS inclusion criteria). These additional data are incorporated into NVDRS. Analysis files are updated monthly at CDC. On the basis of previous experience, CDC estimates that case counts might increase 1%–2% after the initial 18-month data collection period.

Fatal Injuries During 2007

This report provides preliminary data concerning fatal injuries meeting the NVDRS case definition in 2007 for 16 participating states that were received by CDC as of August 31, 2009. Data from California were not included in this report because NVDRS was implemented only in a limited number of cities and counties rather than statewide. Participating states used vital statistics death certificate files to identify deaths meeting NVDRS case definitions. Each state reported all deaths of their residents that occurred within the state and deaths of state residents that occurred elsewhere. Once a death was identified, NVDRS data abstractors linked source documents, linked deaths within each incident, coded data elements, and wrote a short narrative of the incident. These narratives were reviewed for all incidents in which coded data were unclear or incomplete. State-level data then were consolidated and analyzed for this aggregate report. Numbers, percentages, and crude rates are presented in aggregate for all deaths by abstractor-assigned manner of death and for special situations and populations (e.g., homicide followed by suicide, suicides of former or current military personnel, and intimate-partner-related homicides). Rates for cells with a frequency of <20 are not reported because of the instability of those rates. In addition, rates could not be calculated for variables (e.g., marital status, precipitating circumstances) because denominators were unknown. Bridged-race 2007 population estimates were used as denominators in the rate calculations (6). For compatible numerators for rate calculations to be derived, person records listing multiple races were recoded to a single race when possible, using a bridging algorithm provided by NCHS (available at http://www.cdc.gov/nchs/nvss/bridged_race.htm).

Results

All Deaths

Deaths by Manner, Method, and Location

The 16 NVDRS states included in this report collected data concerning 15,882 incidents and 16,319 deaths that occurred during 2007. The crude death rate was 19.9 deaths per 100,000 population. Suicides ($n = 9,245$) accounted for the highest rate of violent death (11.6 per 100,000 population) followed by homicide/legal-intervention ($n = 4,563$) deaths (5.7 per 100,000 population). Deaths of undetermined intent ($n = 2,403$) and unintentional firearm deaths ($n = 107$) occurred at lower rates (3.0 and 0.1 per 100,000 population, respectively). Of all incidents occurring in 2007 in the 16 states included in this report, 2.2% were known to have multiple victims. Firearms accounted for 48.2% of injury deaths, poisoning for 20.4%, and hanging/strangulation/suffocation for 14.2% (rates: 9.9, 4.2, and 2.9 per 100,000 population, respectively); rates for other methods were lower. For all deaths, a house or apartment was the most common location (69.9%). The next-most-common location of injury (8.2%) was a street or highway (Table 1).

Toxicology Results of Decedent

Tests for alcohol were conducted for 73.8% of decedents, and drug tests for amphetamines, antidepressants, cocaine, marijuana, and opiates were conducted for 52.5%, 43.2%, 55.8%, 36.5%, and 54.1% of decedents, respectively. Among decedents who tested positive for alcohol (33.4%), 59.1% had a blood alcohol concentration (BAC) of >0.08 g/dL (the legal limit in the majority of states). Opiates, including heroin and prescription pain killers, were identified in 26.2% of cases tested for these substances (antidepressants [23.5%], cocaine [13.5%], marijuana [11.1%], and amphetamines [4.6%]) (Table 2).

Suicides

Sex, Race/Ethnicity, Age Group, and Marital Status

The 16 NVDRS states included in this report collected data concerning 9,233 fatal suicide incidents and 9,245 suicides that occurred during 2007. Rates of suicide by month showed little variation throughout the year (range: 0.8–1.1 per 100,000 population) (Table 3). Overall, the crude suicide rate was 11.6 per 100,000 population. The rate for males was more than three times that for females (18.4 and 5.0 per 100,000 population, respectively). Non-Hispanic whites accounted for

the largest number of suicide deaths, and AI/ANs and non-Hispanic whites had the highest rates of suicide (18.2 and 14.0 per 100,000 population, respectively). The highest rates of suicide by age group occurred among persons aged 45–54 years, 75–84 years, and 35–44 years (17.6, 16.4, and 16.3 per 100,000 population, respectively). Children aged 10–14 years had the lowest rates of suicide among all age groups (0.8 per 100,000 population). Rates of suicide among adolescents aged 15–19 years (6.9 per 100,000 population) were approximately half of those for persons aged ≥ 30 years (Table 4).

Decedents aged 35–64 years accounted for 55.0% of suicide deaths among males. Rates among males were highest for those aged ≥ 85 years followed by those aged 75–84 years (43.9 and 35.8 per 100,000 population, respectively). AI/AN males had the highest rates of any racial/ethnic population and had rates that were more than four times the rate for API males. Among females, decedents aged 35–64 years accounted for 65.0% of suicides. Rates for females peaked at 9.0 per 100,000 among those aged 45–54 years. As with males, female suicide rates were highest among AI/ANs (7.3) followed closely by non-Hispanic whites (6.2). Among females, the lowest rates of suicide were among non-Hispanic blacks (1.5) and Hispanics (1.7). Of all decedents aged ≥ 18 years for which marital status was known, 38.7% were married, 29.3% had never married, and 22.1% were divorced at the time of death (Table 4).

Method and Location of Injury

Firearms were used in the majority (50.7%) of suicide deaths, followed by hanging/strangulation/suffocation (23.1%) and poisoning (18.8%) (Table 5). The most common method used by male suicide decedents was a firearm (56.0%) followed by hanging/strangulation/suffocation (24.4%). Among females, poisons were used most often (40.8%) followed by firearms (31.9%). The most common place of self-inflicted injury was a house or apartment (77.2%) followed by natural areas (4.4%), and streets or highways (3.0%). A total of 120 (1.3%) suicides occurred in a jail or prison setting (115 males and 5 females) (Table 5).

Toxicology Results of Decedent and Precipitating Circumstances

Tests for alcohol were conducted for 68.9% of suicide decedents, and drug tests for amphetamines, antidepressants, cocaine, marijuana, and opiates were conducted for 44.2%, 39.8%, 47.2%, 35.7%, and 47.1% of suicide decedents, respectively. Among suicide decedents who tested positive for alcohol (34.5%), 62.0% had a BAC of >0.08 g/dL. Opiates (e.g., heroin, prescription pain killers) were identified in 22.8% of cases tested for these substances; cocaine and marijuana were identified in 8.9% and 7.5% of tested cases, respectively. Of

suicide decedents who were tested for antidepressants, 30.0% were positive at the time of their death (Table 6).

Precipitating circumstances were known for approximately 86% of suicide decedents. Overall, mental-health problems were the most commonly noted circumstance for suicide decedents with 41.5% described as experiencing a depressed mood at the time of their deaths. Nearly 45% were described as having a mental-health problem although only 33.8% were receiving treatment (Table 7). Among those with a current mental-health problem, 74.9% had received a diagnosis of depression/dysthymia, 14.5% had been diagnosed with bipolar disorder, and 8.1% with an anxiety disorder (Table 8). Among suicide decedents with known circumstance information, 19.9% had a history of previous suicide attempts, 28.0% disclosed their intent prior to dying, and 33.3% left a suicide note (Table 7). Other than mental health conditions, circumstances noted most often were a crisis in the preceding 2 weeks or intimate-partner problems, each indicated in approximately 30% of suicides with known circumstance information. Physical-health problems also were noted in 21.4% of cases with circumstance information.

Similar percentages of male and female suicide decedents were observed to have a depressed mood at the time of death; however, a higher percentage of females than males had a current mental-health problem (62.9% and 39.6%, respectively) or were being treated for a mental-health problem (50.6% and 28.8%, respectively). Approximately the same percentage of male and female suicide decedents experienced physical-health problems in the period before their deaths, although a higher percentage of males than females had job, financial, or criminal problems in the period preceding their deaths. Intimate-partner problems also were cited as a precipitating factor in a higher percentage of male suicides than female suicides (31.7% and 26.4%, respectively). Although occurring in only a limited percentage of cases, being a perpetrator of interpersonal violence in the month before death was more common among male suicide decedents (5.6%) than being a victim of such violence (0.2%) whereas the proportions were similar for females (1.6% and 1.1%, respectively) (Table 7).

Homicides

Sex, Race/Ethnicity, Age Group, Marital Status

The 16 NVDRS states included in this report collected data concerning 4,324 homicide incidents and 4,563 homicides that occurred during 2007. Overall, the crude homicide rate was 5.7 deaths per 100,000 population in 2007. Rates of homicide by month showed little variation throughout the year (range: 0.4–0.6 per 100,000 population) (Table 9).

The majority (53.0%) of homicide decedents aged ≥ 18 years for which marital status was known had never been married, and 22.7% were married at the time of their death. In 41.2% of homicides, the relation of the victim to the suspect was not known. When a suspect was identified, the suspect most often was an acquaintance or friend (15.7%), a spouse or intimate partner (10.5%), or a stranger (8.4%). Perpetrators were other relatives of the decedent in <10% of cases with known information about the relation of the victim to the suspect (Table 10).

The homicide rate for males was approximately 3.6 times that for females (9.0 and 2.5 per 100,000 population, respectively). Non-Hispanic blacks accounted for the majority (52.0%) of homicide deaths and had the highest rate (19.3 deaths per 100,000 population) followed by AI/ANs (10.5) and Hispanics (7.2). Age-specific homicide rates were highest (14.7 deaths per 100,000 population) among those aged 20–24 years followed by those aged 25–29 years (12.4 deaths per 100,000 population). The rate for infants aged <1 year was more than four times that for children aged 1–4 years (9.7 and 2.2 per 100,000 population, respectively) and similar to that for adolescents aged 15–19 years (9.5 per 100,000 population). Rates were lowest among children aged 5–14 years and persons aged ≥ 65 years. The majority (62.3%) of male homicide decedents were aged 20–44 years; males aged 20–24 years had the highest rates of homicide (24.5 per 100,000 population). For females, homicide rates were highest (8.2 deaths per 100,000 population) among infants aged <1 year (Table 11).

Method and Location of Injury

Firearms were used in 66.1% of homicides, followed by sharp instruments (12.1%) and blunt instruments (5.8%). No other single method was used in more than 3.4% of homicides (Table 9). Firearms were the most common method used in homicides of males (71.7%) and females (46.4%). Hanging/strangulation/suffocation was over six times more common among female homicide decedents than among males (8.6% and 1.4%, respectively). A house or apartment was the most common location of homicide for both males and females (44.7% and 72.6%, respectively). The next-most common location of homicide for males was a street or highway (26.0%), a parking lot or public garage (5.4%), and a motor vehicle (4.0%); for females, the next-most common locations were a street or highway (6.8%), or a commercial/retail or natural area (2.7% each) (Table 12).

Toxicology Results of Decedent and Precipitating Circumstances

Tests for alcohol were conducted for 77.9% of homicide decedents, and drug tests for amphetamines, antidepressants,

cocaine, marijuana, and opiates were conducted for 54.4%, 34.8%, 58.8%, 33.8%, and 52.4% of homicide decedents, respectively. Among homicide decedents who tested positive for alcohol (34.6%), 55.4% had a BAC of >0.08 g/dL. Marijuana, cocaine, and opiates were identified in 20.0%, 15.7%, and 6.8% of homicide decedents tested, respectively (Table 13).

Precipitating circumstances were identified for 64.9% of homicide deaths. Approximately one third of those homicides were precipitated by another crime. In 76.0% of cases precipitated by another crime, the crime was in progress at the time of the incident (Table 14). The crime was most often robbery (37.4%), followed by assault (23.3%), burglary (9.3%), drug-related (8.2%), rape/sexual assault (3.3%), or motor-vehicle theft (3.0%) (Table 15). Other common precipitating circumstances were an argument, abuse, or conflict over something other than money or property (37.5%); drug-related circumstances (13.3%); justifiable self defense (8.0%); or an argument about money or property (5.4%). In 20.4% of cases with known circumstance information, intimate-partner violence was identified as a contributing factor. In approximately 1% of the cases, the decedent was a police officer killed in the line of duty or an intervening person assisting a crime victim (Table 14).

An argument, abuse, or a conflict unrelated to money or property was a factor in more homicides among males than females (41.9% and 24.5%, respectively). Drug-related homicides accounted for 15.0% of male homicides and 8.2% of female homicides. Intimate-partner violence was a precipitating factor in 52.5% of female homicides but only 9.5% of male homicides. In 13.7% of male homicides with known circumstance information, the decedent also used a weapon during the altercation, compared with 1.9% of female homicides (Table 14).

Deaths of Undetermined Intent

Sex, Race/Ethnicity, Age Group, Education, and Marital Status

The 16 NVDRS states included in this report collected data concerning 2,392 incidents involving 2,403 deaths during 2007 for which a determination of intent could not be made. Rates of undetermined death by month were at 0.2 or 0.3 per 100,000 population throughout the year (Table 16). Overall, the crude rate for undetermined deaths was 3.0 per 100,000 population. Rates of undetermined death were higher among males than females (3.8 and 2.3 per 100,000 population, respectively). Although non-Hispanic whites accounted for 71.5% of undetermined deaths, rates were highest among AI/ANs (6.1 per 100,000 population). Nearly half (49.1%) of decedents for whom the manner of death was undetermined

were aged 35–54 years. Rates were highest (22.0 per 100,000 population) among infants aged <1 year. Among decedents aged >18 years with an undetermined manner of death for which marital status was known, 39.3% had never been married, 27.9% were married, and 24.3% were divorced at the time of death. AI/AN males had the highest rates (7.5 per 100,000 population) of undetermined death compared with males or females of any other racial/ethnic population (Table 17).

Method and Location of Injury

The most common method of injury was poisoning (65.0%). No other known single method accounted for >2.7% of undetermined deaths. Among both males and females for which the method of injury was known, poisoning was reported for 63.6% and 67.4% of deaths, respectively. The majority of undetermined deaths occurred in a house or apartment, making it the most common place of injury for both males and females (75.1% and 82.6%, respectively). A natural area was the second most common setting, accounting for 3.6% of deaths among males and 2.5% among females (Table 18).

Toxicology Results of Decedent and Precipitating Circumstances

Tests for alcohol were conducted for 85.4% of decedents of undetermined intent, and drug tests for amphetamines, antidepressants, cocaine, marijuana, and opiates were conducted for 80.8%, 72.4%, 83.4%, 44.8%, and 84.9% of decedents, respectively. Among decedents who tested positive for alcohol (28.2%), 55.2% had a BAC of >0.08 g/dL. Among decedents tested for opiates, 56.6% were positive; of those tested for cocaine, 21.1% were positive; of those tested for marijuana, 9.4% were positive; and of those tested for antidepressants, 27.8% were positive (Table 19).

Precipitating circumstances were known in approximately 73% of deaths of undetermined intent. Of those, 28.7% of decedents had a problem with alcohol, and 62.4% had other substance-abuse problems (e.g., illicit drug or prescription abuse). Although a current depressed mood was reported for only 13.0% of decedents, 39.0% of decedents with known circumstance information had a current mental-health problem, 31.7% were receiving treatment at the time of their death, 10.5% had a history of suicide attempts, 7.2% had disclosed an intent to commit suicide, and 1.6% left a suicide note. Other circumstances noted most often were physical-health problems (29.0%), a crisis during the preceding 2 weeks (15.0%), or an intimate-partner problem (10.8%) (Table 20). Of those with a current mental-health problem, 61.2% had received a diagnosis of depression/dysthymia, 16.0% of bipolar disorder, and 12.5% of an anxiety disorder (Table 21).

A greater percentage of male than female decedents were reported to have an alcohol problem (33.1% and 21.6%, respectively) or other substance-abuse problems (65.1% and 57.9%, respectively) at the time of death. Mental-health problems were reported in a higher percentage of undetermined deaths of females than of males (52.0% and 30.9%, respectively), and a higher percentage of females were receiving treatment for a mental-health problem than males (44.6% and 23.7%, respectively) and had a history of suicide attempts (13.8% and 8.4%, respectively) (Table 20).

Unintentional Firearm Deaths

Sex, Race/Ethnicity, Age Group, and Seasonality

The 16 NVDRS states included in this report collected data concerning 107 unintentional firearm deaths during 2007. Males accounted for 90.7% of decedents. The majority (68.2%) were non-Hispanic whites, followed by non-Hispanic blacks (22.4%). Approximately 20% of unintentional firearm fatalities occurred among persons aged 15–19 years. November had the highest percentage of unintentional firearm deaths (17.8%) at nearly twice the frequency of any other month (Table 22).

Location of Injury

Approximately 64.5% of all unintentional firearm fatalities took place in a house or apartment, making it the most common place of injury for both males and females, followed by natural areas (16.8%) (Table 22).

Context of the Injury and Associated Circumstances

Overall, unintentional firearm injury deaths occurred more commonly while victims were playing with a gun (29.9%), hunting (24.7%), showing a gun to others (14.3%), or loading or unloading a gun (10.4%). The circumstances of injury included thinking that a gun was unloaded, unintentionally pulling the trigger, and experiencing a gun malfunction (26.0%, 19.5%, and 5.2%, respectively) (Table 23).

Special Topics

Violent Deaths with Multiple Decedents

The 16 NVDRS states included in this report collected data on 347 incidents that resulted in multiple decedents. Firearms were the most common method (73.7%) used in incidents with multiple decedents, followed by poisonings (5.7%), sharp instruments or hanging/strangulation/suffocation (5.2% each), and blunt instruments (3.3%) (Table 24). Of a total of

784 victims, 461 (58.8%) were males; 334 (90.8%) of 368 suspects also were males. Non-Hispanic whites accounted for the highest percentage of decedents (57.0%), followed by non-Hispanic blacks (26.5%) and Hispanics (10.5%). Rates for decedents were highest for non-Hispanic blacks and persons aged 15–54 years. Suspects most commonly were aged 20–54 years (Table 25).

Homicide Followed by Suicide

The 16 NVDRS states included in this report collected data concerning 172 violent incidents that occurred during 2007 in which a homicide was followed by the suicide of the suspect. Of 240 homicide decedents, 174 (72.5%) were female and 160 (93.0%) suspects (suicide decedents) were male. Homicide rates were similar (0.3 per 100,000) among racial/ethnic groups which had results reported; 70.4% of homicide decedents were non-Hispanic whites. Among suspects who killed themselves after committing homicide, 66.9% were non-Hispanic whites and 15.7% were non-Hispanic blacks. The highest percentages of both homicide and suicide decedents were aged 35–54 years (31.7% and 49.4%, respectively) (Table 26).

The majority of homicide decedents and suspects (34.7% and 32.4%, respectively) were married at the time of death (not necessarily to each other) (Table 26). With respect to location, 75.4% of the homicides occurred in a house or apartment and 2.1% each in a street/highway or commercial/retail area. Firearms were the most common (approximately 80%) method used by suspects both in committing the homicide and in subsequently killing themselves (Table 27).

Tests for alcohol were conducted for 79.6% of homicide decedents and 72.7% of suicide decedents. Among decedents who tested positive for alcohol (18.3% of homicide victims; 32.8% of suicide decedents), 48.6% of homicide decedents and 53.7% of suicide decedents had a BAC of >0.08 g/dL at the time of death. Suspects who killed themselves following a homicide and who were tested subsequently for drugs had higher percentages of positive tests for antidepressants, cocaine, marijuana, and opiates than homicide victims (Table 28).

Although 8.3% of persons who killed themselves following a homicide had a current depressed mood, only 3.6% were receiving mental-health treatment at the time of the fatal incident. Intimate-partner-relationship problems preceded homicide followed by suicide in 81.0% of suspected suicides. Other nonintimate-partner-relationship problems contributed to 13.7% of suspected suicides. Of suspects who killed themselves, 91.1% had had a personal crisis within the preceding 2 weeks. Previous criminal legal problems were noted in 19.1% of suspected suicides and noncriminal problems in 3.0%; physical health or financial problems were contributing circumstances in 6.6% and 4.2% of suspected suicides, respectively; 6.0% of

suicide decedents had disclosed their intent to kill themselves; and 1.8% had a history of suicide attempts (Table 29).

Intimate-Partner Homicide

The 16 NVDRS states included in this report collected data concerning 562 incidents comprising 612 deaths of intimate-partner-related homicides that occurred during 2007. Of 612 homicide victims, 394 (64.4%) were female. Although 51.3% of homicide victims were non-Hispanic whites, rates were higher for AI/ANs and non-Hispanic blacks (2.1 and 1.6 per 100,000 population, respectively). Of 580 suspects, 451 (77.8%) were male, 252 (43.5%) were non-Hispanic whites and 186 (32.1%) non-Hispanic blacks. The highest percentages of victims and suspects (26.1% and 23.5%, respectively) were persons aged 35–44 years. The highest percentage (37.8%) of victims were married at the time of death (Table 30). Tests for alcohol were conducted for 81.5% of the victims. Of the 33.9% of decedents who tested positive for alcohol, 60.4% had a BAC of >0.08 g/dL. The percentage of victims tested for substances other than alcohol varied (range: 37.8%–56.5%) for various drugs; cocaine was evident in approximately 13% of victims tested for this substance (Table 31).

Suicide of Former or Current Military Personnel

The 16 NVDRS states included in this report collected data concerning 1,774 suicides by former or current military personnel that occurred during 2007. Of these decedents, 1,713 (96.6%) were male, and 1,627 (91.7%) were non-Hispanic whites. The greatest percentage of decedents were persons aged ≥35 years. The most common method (67.2%) used was a firearm followed by hanging/strangulation/suffocation (14.5%) and poisoning (12.3%) (Table 32). Among the 64.2% former or current military personnel suicide decedents who were tested for alcohol, 31.6% tested positive; 64.2% of these decedents had a BAC of >0.08 g/dL (Table 33). Although 43.2% were depressed at the time of death, and 36.8% had a mental-health problem, only 27.4% were receiving mental-health treatment. With respect to substance abuse, 16.4% had an alcohol problem, and 7.9% had a problem with other substances. Among those with known circumstance information, 25.2% had experienced a problem with an intimate partner, 37.2% had a physical-health problem, and 32.4% had experienced an acute crisis during the preceding 2 weeks. With respect to life stressors, 11.0% had experienced a job problem, 10.0% a financial problem, and 7.5% a criminal legal problem. Approximately one third (34.7%) left a suicide note, 13.8% had made a previous suicide attempt, and 26.9% had disclosed an intent to commit suicide (Table 34).

Legal Intervention

The 16 NVDRS states included in this report collected data on 145 legal-intervention incidents in 2007 resulting in 141 single-victim deaths and four deaths where the legal-intervention victim had recently committed a homicide. Of the 145 legal-intervention decedents, 45.5% were non-Hispanic whites and 35.2% were non-Hispanic blacks. With respect to location, 37.9% of legal-intervention deaths occurred in a house or apartment, 31.7% on a street or highway, and 9.0% in a parking lot or public garage (Table 35). The majority of decedents were aged 20–54 years (Table 36). Of the 87.6% of legal-intervention decedents tested for alcohol, 38.6% were positive for alcohol and 67.4% of these decedents had a BAC of >0.08 g/dL. The percentage of victims tested for other substances varied (range: 49.7%–75.9%). The presence of other drugs for which tests were positive also varied: 30.6% of those tested for marijuana, 24.6% of decedents tested for cocaine, 20.4% of those tested for amphetamines, 9.3% of those tested for opiates, and 5.4% of those tested for antidepressants were positive for these substances (Table 37).

Suicide Among Persons Aged ≥50 Years

In 2007, NVDRS collected data for 3,677 persons aged ≥50 years who died by suicide. Of those, rates of suicide were highest among those aged 50–59 years (16.7 per 100,000 population), followed by those aged ≥80 years (15.8 per 100,000). Beginning with those aged 60–69 years, rates increase with age from 13.4 per 100,000 to 14.4 for those aged 70–79 years, and 15.8 for those aged ≥80 years. Among persons aged ≥50 years, rates were five times higher among males than among females (26.1 and 5.1 per 100,000 population, respectively). Rates were highest among non-Hispanic whites (17.7 per 100,000 population), followed by AI/ANs (11.7 per 100,000 population), Hispanics (8.6), APIs (5.6), and non-Hispanic blacks (3.9). At the time of death, persons aged 50–69 years most often were either married or divorced. Those aged 70–79 years and those aged ≥80 years most often were either married or widowed (Table 38).

The majority (81.9%) of suicide decedents aged ≥50 years died in a house or apartment. The second-most-common location for those aged 50–59 and 60–69 years was a natural area (4.7% and 3.5%, respectively). The second-most-common location for those aged 70–79 years was a street/highway (2.2%) and for those aged ≥80 natural areas, street/highway and motor vehicle were equally common at 1.2% each. As to method used by suicide decedents aged ≥50 years, firearms accounted for 60.6% of deaths (rate: 9.3 per 100,000 population), poisoning for 19.6% (3.0 per 100,000 population), and hanging/strangulation/suffocation for 12.6% (1.9 per 100,000

population). Rates of firearm suicide were highest among persons aged ≥80 years (12.1 per 100,000) and those aged 70–79 years (11.1 per 100,000 population) (Table 38).

Precipitating circumstances were identified for approximately 86% of older adult suicides. Current depressed mood (39.1%), current mental-health problem (39.1%), and physical-health problems (33.0%) were the most commonly identified circumstances; 30.8% left a suicide note, and 24.7% disclosed their intent to commit suicide (Table 39).

Discussion

The findings in this report indicate clear variations in patterns of death from violence-related injuries, unintentional firearm-related injuries, and deaths of undetermined intent reported from the 16 states included in this report. Rates for these deaths were disproportionately higher among males, adults aged <55 years, and minority populations. A residence (house or apartment) was the most common location for all deaths. Of all incidents meeting NVDRS inclusionary criteria in 2007 in the 16 states included in this report, approximately 98% involved a single victim.

Suicide Patterns

Suicide rates were higher among males than among females, AI/ANs, and non-Hispanic whites than among non-Hispanic blacks, and highest among persons aged 45–54 years. These findings are similar to those that have been documented in other reports (7–9). Persons aged ≥80 years have typically had the highest rates of suicide in the United States (1). However, in 2006, rates of suicide among persons aged 45–54 years in the United States surpassed those for persons aged ≥80 years (1). This pattern also was noted in the findings from NVDRS states in 2006 (10) and has continued in 2007. Problems related to mental health, jobs, finances, or relationships might have contributed to the high rates of suicide in this age group. Mental health and/or substance-abuse problems, relationship problems and losses, and recent crises were frequent precipitants for suicide. These factors have been documented in other studies as important risk factors for suicide (8,11).

Alcohol was a factor in approximately one third of the reported suicides, and 62% of these decedents had a BAC of >0.08 g/dL at the time of death. Alcohol and drug abuse are second only to depression and other mood disorders as the most frequent risk factors for suicidal behavior (8,12,13). However, the relation between these factors is complex. Previous research indicates that alcohol intoxication might act as an important proximal (i.e., an experience often occurs immediately prior to a suicide) risk factor for suicidal behavior (14). In cases

where there is alcohol dependency, the alcohol abuse might lead directly to depression or indirectly through a sense of decline or failure that is experienced by many persons who are dependent on alcohol. Alcohol also might be a form of self-medication to alleviate depression. Both depression and alcohol abuse also might be the result of specific stresses in a person's life (15). The co-occurrence of mood disorders with substance abuse, including alcohol abuse, greatly increases the risk for suicidal behavior (16).

Many of the circumstances surrounding suicides (e.g., mental health problems, alcohol problems, and recent crises) also were found for certain special populations examined in this report. However, other factors also were important. Physical health problems were more frequently noted for former or current military personnel and in adults aged ≥ 70 years. Physical illness is believed to be a common antecedent to suicide among the elderly, although prevalence estimates vary widely, and it appears to be a stronger contributing factor when mood disorders, depressive symptoms or other factors also are present (8,15). For example, untreated or undertreated pain, anxiety about the progression of an illness, fear of dependence, and fear of burdening family members are major contributing factors in suicidal behavior among elderly persons with physical illnesses (8). Many of these factors also are evident in military populations with disabling conditions. The proportion of former or current military personnel reported to be experiencing health problems also might reflect a difference in reporting and contact with health-care professionals.

Finally, approximately 30% of suicide victims had disclosed their intent to commit suicide, and approximately 20% had made a previous suicide attempt. A previous suicide attempt is an important predictor of subsequent fatal suicidal behavior (8,15). Disclosure of intent also is an important warning sign of suicidal intentions, although persons in close contact with potential victims of suicide often are unaware of the significance of these warnings or unsure how to act on them (17).

Homicide Patterns

Homicide rates were higher among males than among females, among non-Hispanic blacks compared with members of other racial/ethnic populations, and among persons aged 20–24 years compared with persons in other age groups. These findings also are consistent with patterns documented in other reports. Homicide is the second leading cause of death in the United States among persons aged 15–24 years, and rates among non-Hispanic blacks in this age group exceed those of other racial/ethnic populations by approximately fourfold to sevenfold (1). Males also are disproportionately represented among victims of homicide in the United States and elsewhere (1,18).

The majority of homicides involved a single victim. Multiple decedent homicides and homicide-suicide incidents accounted for <3% of violent deaths. The majority of homicides were related to interpersonal conflicts. Crime was a factor in approximately one third of all homicide/legal-intervention deaths, with robbery being the primary circumstance. These findings are consistent with other research on homicide. Arguments and conflicts are immediate motivations for the majority of both male and female homicides in the United States (19). One factor that distinguishes male from female homicides is the relationship between the victim and the perpetrator. In the United States, approximately one in three homicides of females is committed by a current or former spouse or partner (20). Among male homicide victims, approximately 5% are killed by intimate partners. The findings of this report indicate that male homicide decedents were more frequently killed following arguments or conflicts with persons other than an intimate partner or for other reasons (e.g., crime or drug-related) whereas more than half of homicides involving a female victim involved intimate-partner-related violence.

As with suicide decedents, alcohol was present in approximately one third of homicide decedents; more than half of these decedents (55.4%) had BACs of >0.08 g/dL. Alcohol is an important situational factor in interpersonal violence. In the case of interpersonal violence among youths, excessive alcohol consumption might increase impulsivity and make some drinkers more likely to resort to violence in a confrontation or argument (21,22). Reduced physical control and the ability to assess risks in potentially dangerous situations also can make some drinkers more vulnerable to victimization (21,22). In the case of intimate-partner violence, excessive alcohol consumption by one or both partners might exacerbate financial or child care problems or other stressors and increase tension and conflict in the relationship (23). Alcohol also can be a form of self-medication to cope with previous or current experiences of abuse (23).

Method of Injury

Approximately two thirds of all homicides and approximately one half of all suicides in the United States are committed with a firearm (1). In the 16 states included in this report, firearms were the most common method used in homicides, incidents involving multiple victims, and incidents of homicide followed by suicide. Previous research indicates that interpersonal disputes can escalate and cause serious violent injury or death, especially when weapons of lethal means (e.g., firearms) are involved in the dispute (24,25). Firearms also were the most common method used in suicides; however, methods differed by sex. Firearms were the most common method used by

males to complete suicide followed by hanging/strangulation/suffocation. Poisoning was the most common method used by females, followed by firearms.

Deaths where the intent was undetermined were primarily the result of poisonings or had an unknown cause. Poisoning was the most common method for both males and females. Toxicology results documented a high prevalence of alcohol and other substances at the time of death. For example, approximately 85% of decedents with undetermined intent were tested for opiates, and nearly 57% tested positive for these substances. Whether these deaths were related to unintentional drug poisonings, (which have increased substantially in recent years, particularly among adults aged 35–54 years [26,27]) or were suicides is unknown. The majority (49%) of decedents in the 16 states were aged 35–54 years. Substance-abuse problems involving drugs other than alcohol were the most commonly noted circumstance; approximately 40% of such decedents had a mental-health problem, and 10% had a history of suicide attempts.

Prevention Opportunities

Information concerning the precipitating circumstances in violent deaths described in this report provides important clues regarding where to focus prevention efforts. For example, relationship problems, interpersonal conflicts, and recent crises were important precipitating factors for both homicide and suicide. Intimate-partner-related problems, in particular, were a factor in many types of violent death. Primary prevention programs designed to enhance social problem-solving and coping skills to deal with stressful life events, health and financial problems, or other problems that occur within interpersonal relationships can potentially reduce violence (28). In addition to demonstrating the need to address situational stressors, the findings in this report underscore the importance of changing cultural and social norms (e.g., attitudes condoning the use of violence as a means of resolving conflict) addressing the social and economic conditions within communities that often give rise to violence (e.g., inequities with regard to the distribution of and access to resources and opportunities, social isolation, lack of connectedness among persons, families, and communities) and intervening much earlier by teaching young persons the skills to develop and promote respectful, nonviolent interpersonal relationships. Some of the strategies that offer the strongest evidence of effectiveness with respect to the latter are primary prevention strategies that focus on family environments, school environments, and building individual social, emotional, and behavioral competencies (28–31).

The findings of this report also highlight the importance of addressing mental health problems. Mental-health problems

were highly prevalent among suicide decedents, yet many were not receiving treatment at the time of death. Reasons why persons do not seek care for mental health problems include the belief that the problem will resolve on its own, financial barriers, a lack of awareness of available services, fear of hospitalization, embarrassment, fear of what others might think, and the belief that acknowledgment or discovery of a mental health problem could damage one's career or relationships (8). Despite public education efforts focused on the nature, causes, and treatment of mental illness, the stigma of mental illness is one of the most significant barriers deterring persons from seeking treatment (8). Stigma leads persons to fear, reject, and distance themselves from persons with mental health problems. Findings from a national study indicate that approximately 47%–63% of persons in the United States want to distance themselves from persons with depression and schizophrenia (32). The consequences of stigma for persons suffering from these problems include diminished opportunities, lowered self-esteem, shame and concealment of symptoms, and lower help-seeking behavior (8). These results underscore the need for prevention measures aimed at changing cultural attitudes and norms surrounding mental health problems so that persons in need of treatment can seek the care of professionals and reach out to family, friends, and others without hesitation.

Limitations

The findings provided in this report are subject to at least seven limitations. First, the availability, completeness, and timeliness of data are dependent on the sharing of data among state health department NVDRS teams, CMEs, and law enforcement personnel in their states. This is particularly challenging when states have independent county coroner systems rather than a centralized CME system and large numbers of law enforcement jurisdictions. NVDRS incident data might be limited or incomplete for areas in which these data-sharing relations are not developed fully.

Second, toxicology data are not collected consistently across all states or for all alcohol and drug categories. The percentage of decedents testing positive might be affected by selective testing biases in medical examiner or coroner offices (33). Third, abstractors are limited to the data included in the reports they receive. Reports might not fully reflect all information known about an incident, particularly in the case of homicides, when data are less readily available until after prosecutions are complete. Fourth, case definitions present challenges when a single death is classified differently in different documents (e.g., “unintentional” in a police report, “homicide” in a CME report, and “undetermined” on the death certificate). NVDRS abstractors reconcile these cases using standardized NVDRS

case definitions and select a single manner of death on the basis of all source documents. Fifth, NVDRS data are available only from a limited number of states and therefore are not nationally representative. Sixth, although extensive coding training is conducted and help desk support is available daily, variations in coding might occur depending on the abstractor's level of experience. For this reason, states regularly conduct blinded reabstraction of cases to test consistency and identify training needs. Finally, protective factor data (i.e., characteristics or circumstances that reduce the risk for violent death) are not collected by NVDRS because of the nature of death certificate, CME record, and police reports, which typically contain only circumstances associated with risk factors.

Conclusion

Accurate, timely, and comprehensive surveillance data can be used to monitor the occurrence of violence-related fatal injuries and assist public health and other authorities in the development, implementation, and evaluation of programs and policies that reduce and prevent violent deaths and injuries at the national, state, and local levels (34,35). Continued development and expansion of NVDRS is critical to the public health and criminal justice communities at the federal, state, and local levels that work to reduce the personal, familial, and societal costs of violence. Further efforts are needed to increase the number of states participating in NVDRS, with the ultimate goal of full national representation, including all 50 states, the District of Columbia, and U.S. territories.

Acknowledgments

Contributors to this report included participating state Violent Death Reporting Systems; participating state agencies, including state health departments, vital registrars' offices, coroners' and medical examiners' offices, crime laboratories, and local and state law enforcement agencies; partner organizations, including the State and Territorial Injury Prevention Directors' Association, National Violence Prevention Network, National Association of Medical Examiners, National Association for Public Health Statistics and Information Systems, Council of State and Territorial Epidemiologists, the International Association of Chiefs of Police, and Association of State and Territorial Health Officials; federal agencies, including the Department of Justice (Bureau of Justice Statistics and the Federal Bureau of Investigation), the Department of the Treasury (Bureau of Alcohol, Tobacco, and Firearms); other stakeholders, researchers, and foundations, including Harvard University School of Public Health, the Joyce Foundation, and Fenton Communications; and the National Institute for Occupational Safety and Health, National Center for Health Statistics, CDC.

References

1. CDC. Web-based Injury Statistics Query and Reporting System (WISQARS™). Atlanta, GA: US Department of Health and Human Services, CDC; 2008. Available at <http://www.cdc.gov/ncipc/wisqars/default.htm>.
2. Doll L, Bonzo S, Mercy J, et al, eds. Handbook of injury and violence. New York, NY: Springer; 2007.
3. Paulozzi LJ, Mercy J, Frazier L, et al. CDC's National Violent Death Reporting System: background and methodology. *Inj Prev* 2004;10:47–52.
4. CDC. Surveillance for violent death—National Violent Death Reporting System, 16 states, 2005. In: CDC Surveillance Summaries, April 11, 2008. *MMWR* 2008;57(No. SS-3).
5. World Health Organization. International classification of diseases, version 10. Geneva, Switzerland: World Health Organization; 2007. Available at <http://www.who.int/classifications/icd/en/index.html>.
6. CDC. U.S. census populations with bridged race categories. Hyattsville, MD: US Department of Health and Human Services, CDC; 2007. Available at <http://www.cdc.gov/nchs/about/major/dvs/popbridge/popbridge.htm>.
7. Kung HC, Hoyert DL, Xu JQ, et al. Deaths: final data for 2005. *Nat Vital Stat Rep* 2008;56:1–124.
8. Institute of Medicine. Reducing suicide: a national imperative. Washington, DC: National Academies of Science; 2002.
9. CDC. Surveillance for fatal and nonfatal injuries—United States, 2001. In: CDC Surveillance Summaries, September 3, 2004. *MMWR* 2004;53(No. SS-7).
10. Karch DL, Dahlberg LL, Patel N, et al. Surveillance for violent deaths—National Violent Death Reporting System, 16 States, 2006. *CDC Surveillance Summaries*, March 20, 2009, *MMWR* 2009;58(No. SS-1).
11. US Department of Health and Human Services. National strategy for suicide prevention: goals and objectives for action. Rockville, MD: United States Department of Health and Human Services, Public Health Service; 2001.
12. Borges G, Walters EE, Kessler RC. Associations of substance use, abuse, and dependence with subsequent suicidal behavior. *Am J Epidemiol* 2000;15:781–9.
13. Tondo L, Baldessarini RJ, Hennen J, et al. Suicide attempts in major affective disorder patients with comorbid substance use disorders. *J Clin Psychiatry* 1999;60(Suppl 2):S63–9.
14. Hufford MR. Alcohol and suicidal behavior. *Clin Psychol Rev* 2001;21:797–811.
15. DeLeo D, Bertolote J, Lester D. Self-directed violence. In: Krug EG, Dahlberg LL, Mercy JA, et al., eds. World report on violence and health. Geneva, Switzerland: World Health Organization; 2002:183–212.
16. Moscicki EK. Epidemiology of completed and attempted suicide: Toward a framework for prevention. *Clinical Neuroscience Research* 2001;1:310–23.
17. American Association of Suicidology. Understanding and helping the suicidal individual. Fact sheet. Available at http://www.suicidology.org/c/document_library/get_file?folderId=232&name=DLFE-30.pdf.
18. Krug EG, Dahlberg LL, Mercy JA, et al. World report on violence and health. Geneva, Switzerland: World Health Organization; 2002.
19. Schwartz, J. Gender differences in homicide offending. In: DeLisi M, Conis P, eds. Violent offenders: theory, research, public policy, and practice, Boston, MA: Jones & Bartlett; 2007:119–40.
20. Federal Bureau of Investigation. Crime in the United States, 2007. Washington, DC: US Department of Justice, Federal Bureau of Investigation; 2007. Available at http://www.fbi.gov/ucr/cius2007/offenses/expanded_information/homicide.html.
21. Parker RN. Alcohol and violence: connections, evidence and possibilities for prevention. *J Psychoactive Drugs* 2004 (Supp2):S157–63.
22. Graham K. Social drinking and aggression. In Mattson MR, ed. Neurobiology of aggression: understanding and preventing violence. 1st ed. Totowa, NJ: Humana Press; 2003:253–74.
23. World Health Organization. Alcohol and intimate partner violence. Fact sheet, 2005. Geneva, Switzerland: World Health Organization; 2005. Available at http://www.who.int/violence_injury_prevention/violence/world_report/factsheets/ft_intimate.pdf.

24. Fagan J, Wilkinson DL. Guns, youth violence, and social identity in inner cities. *Crime and Justice* 1998;24:105–88.
25. Wilkinson DL. Guns, violence and identity among African-American and Latino youth. New York, NY: LFB Scholarly Publishing; 2003.
26. Paulozzi LJ, Budnitz DS, Xi Y. Increasing deaths from opioid analgesics in the United States. *Pharmacoepidemiol Drug Saf* 2006;15:618–27.
27. CDC. Unintentional poisoning deaths—United States, 1999–2004. *MMWR* 2007;56:93–6.
28. Lutzker JR, ed. Preventing violence: research and evidence-based intervention strategies. Washington, DC: American Psychological Association; 2006.
29. Hahn R, Fuqua-Whitley D, Lowry J, et al. The effectiveness of universal school-based programs for the prevention of violence: a report on recommendations of the Task Force on Community Preventive Services. *Am J Prev Med* 2008;33:S114–29.
30. Wilson SJ, Lipsey MW, Derzon JH. The effects of school-based intervention programs on aggressive behavior: a meta-analysis. *J Consult Clin Psychol* 2003;71:136–49.
31. Henggeler SW, Clingempeel WG, Brondino MJ, et al. Four-year follow-up of multisystemic therapy with substance-abusing and substance-dependent juvenile offenders. *J Am Acad Child Adolesc Psychiatry* 2002;41:868–74.
32. Link BG, Phelan JC, Bresnahan M, et al. Public conceptions of mental illness: Labels, causes, dangerousness, and social distance. *Am J Public Health* 1999;89:1328–33.
33. CDC. Toxicology testing and results for suicide victims—13 states, 2004. *MMWR* 2006;55:1245–8.
34. Karch, D, Logan, J. Data consistency in multiple source document: findings from homicide incidents in the National Violent Death Reporting System, 2003–2004. *Homicide Studies*, 2008;12:264–76.
35. Logan J, Karch D, Crosby A. Reducing unknown data in violent death surveillance: a study of death certificates, coroner/medical examiner and police reports from the National Violent Death Reporting System, 2003–2005. *Homicide Studies* 2009;13:385–97.

TABLE 1. Number*, percentage,† and rate§ of incidents, by incident type, manner of death, method used, and location in which injury occurred — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	No.	(%)	Rate
Incident type			
Suicide, single	9,048	(57.0)	NR**
Homicide, single	3,858	(24.3)	NR**
Unintentional firearm	107	(0.7)	NR**
Suicide, multiple	12	(0.1)	NR**
Homicide, multiple	148	(1.0)	NR**
Legal intervention	141	(0.9)	NR**
Homicide followed by suicide	172	(1.1)	NR**
Undetermined	2,389	(15.0)	NR**
Other combinations of deaths	6	—	NR**
Unknown	1	—	NR**
Total	15,882	(100.0)	NR**
Manner of death			
Homicide/Legal intervention	4,563	(28.0)	5.7
Suicide	9,245	(56.6)	11.6
Undetermined intent	2,403	(14.7)	3.0
Unintentional firearm	107	(0.7)	0.1
Unknown	1	—	NR††
Total	16,319	(100.0)	19.9
Method			
Firearm	7,871	(48.2)	9.9
Sharp instrument	712	(4.4)	0.9
Blunt instrument	321	(2.0)	0.4
Poisoning	3,328	(20.4)	4.2
Hanging/Strangulation/Suffocation	2,312	(14.2)	2.9
Personal weapons (hands, feet, fists)	161	(1.0)	0.2
Fall	205	(1.3)	0.3
Drowning	156	(1.0)	0.2
Fire/Burns	77	(0.5)	0.1
Shaking	28	(0.2)	—
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	169	(1.0)	0.2
Intentional neglect	20	(0.1)	—
Other (single method)	138	(0.8)	0.2
Firearm and poisoning§§	1	—	NR††
Firearm and other method type§§	35	(0.2)	—
Poisoning and other method type§§	56	(0.3)	0.1
Other combination of methods††	107	(0.7)	0.1
Unknown	622	(3.8)	0.8
Total	16,319	(100.0)	19.9
Location			
House	11,406	(69.9)	14.3
Street/Highway	1,344	(8.2)	1.7
Motor vehicle	392	(2.4)	0.5
Bar/Nightclub	69	(0.4)	0.1
Commercial/Retail area	188	(1.2)	0.2
Industrial or construction area	50	(0.3)	0.1
Office building	49	(0.3)	0.1
Parking lot/Public garage	337	(2.1)	0.4
Abandoned house/Building/Warehouse	23	(0.1)	—
Park, playground, sports/Athletic area	255	(1.6)	0.3
Preschool/School/College/School bus	56	(0.3)	0.1

TABLE 1. (Continued) Number*, percentage,† and rate§ of incidents, by incident type, manner of death, method used, and location in which injury occurred — National Violent Death Reporting System, 16 states,¶ 2007

Location	No.	(%)	Rate
Public transportation/Station/Railroad tracks	56	(0.3)	0.1
Hospital or medical facility	68	(0.4)	0.1
Supervised residential facility	94	(0.6)	0.1
Jail/Prison	153	(0.9)	0.2
Farm	48	(0.3)	0.1
Natural area	627	(3.8)	0.8
Hotel/Motel	234	(1.4)	0.3
Other	479	(2.9)	0.6
Unknown	391	(2.4)	0.5
Total	16,319	(100.0)	19.9

* No. victims = 16,319 (79.4%); no. suspects/victims = 194 (0.9%); no. live suspects = 4,202 (20.4%); no. persons with unknown role = 29 (0.1%); no. incidents = 16,319.

† Percentages might not total 100% because of rounding.

§ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Because the number of victims varies in incidents involving multiple deaths, population denominators cannot be determined to compute rates.

†† Rates not reported when number of decedents is <20.

§§ Deaths involving more than one method and for which evidence did not indicate which method caused injury.

TABLE 2. Number* and percentage of victims who were tested for alcohol and drugs whose results were positive, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology Variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)§	12,042	(73.8)	4,022	(33.4)
BAC ≤0.08 g/dL§			1,384	(34.4)
BAC >0.08 g/dL§			2,375	(59.1)
Alcohol positive, level unknown			263	(6.5)
Amphetamines	8,559	(52.5)	391	(4.6)
Antidepressants	7,044	(43.2)	1,656	(23.5)
Cocaine	9,103	(55.8)	1,233	(13.5)
Marijuana	5,958	(36.5)	664	(11.1)
Opiates	8,834	(54.1)	2,316	(26.2)
Other Drug(s)	7,911	(48.5)	3,529	(44.6)

* N=16,319.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ BAC of 0.08% g/dL used as the standard for intoxication. Other substances indicated if any results were positive; levels for these substances are not measured.

TABLE 3. Number*, percentage†, and rate§ of suicides, by method used and month in which suicide occurred — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	No.	(%)	Rate
Method			
Firearm	4,683	(50.7)	5.9
Sharp instrument	155	(1.7)	0.2
Blunt instrument	6	(0.1)	NR**
Poisoning	1,742	(18.8)	2.2
Hanging/Strangulation/Suffocation	2,132	(23.1)	2.7
Fall	148	(1.6)	0.2
Drowning	96	(1.0)	0.1
Fire/Burns	30	(0.3)	0.0
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	104	(1.1)	0.1
Other (single method)	20	(0.2)	0.0
Firearm and poisoning††	1	—	NR**
Firearm and other method type††	5	(0.1)	NR**
Poisoning and other method type††	28	(0.3)	0.0
Other combination of methods††	12	(0.1)	NR**
Unknown	83	(0.9)	0.1
Total	9,245	(100.0)	11.6
Month			
January	707	(7.6)	0.9
February	674	(7.3)	0.8
March	799	(8.6)	1.0
April	745	(8.1)	0.9
May	832	(9.0)	1.0
June	773	(8.4)	1.0
July	795	(8.6)	1.0
August	769	(8.3)	1.0
September	796	(8.6)	1.0
October	852	(9.2)	1.1
November	770	(8.3)	1.0
December	719	(7.8)	0.9
Unknown	14	(0.2)	NR**
Total	9,245	(100.0)	11.6

* No. incidents = 9,233; no. victims = 9,245; no. decedents = 9,245.

† Percentages might not total 100% because of rounding.

§ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rates not reported when number of decedents is <20.

†† Deaths involving more than one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 4. Number, percentage*, and rate† of suicides, by decedent's sex, age group, race/ethnicity, and marital status — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	Male			Female			Total		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Age group (yrs)									
10–14	35	(0.5)	1.3	10	(0.5)	NR¶	45	(0.5)	0.8
15–19	324	(4.5)	11.3	61	(3.0)	2.2	385	(4.2)	6.9
20–24	587	(8.1)	20.5	124	(6.1)	4.7	711	(7.7)	12.9
25–29	571	(7.9)	20.0	142	(7.0)	5.2	713	(7.7)	12.7
30–34	575	(8.0)	21.9	146	(7.2)	5.6	721	(7.8)	13.8
35–44	1,446	(20.1)	24.9	460	(22.6)	7.9	1,906	(20.6)	16.3
45–54	1,528	(21.2)	26.5	542	(26.6)	9.0	2,070	(22.4)	17.6
55–64	991	(13.7)	23.4	322	(15.8)	7.1	1,313	(14.2)	14.9
65–74	503	(7.0)	21.8	137	(6.7)	5.0	640	(6.9)	12.7
75–84	467	(6.5)	35.8	66	(3.2)	3.4	533	(5.8)	16.4
≥85	182	(2.5)	43.9	23	(1.1)	2.5	205	(2.2)	15.3
Unknown	2	—	NR¶	1	—	NR¶	3	—	NR¶
Total	7,211	(100.0)	18.4	2,034	(100.0)	5.0	9,245	(100.0)	11.6
Race/ethnicity									
White, non-Hispanic	6,074	(84.2)	22.1	1782	(87.6)	6.2	7,856	(85.0)	14.0
Black, non-Hispanic	484	(6.7)	8.3	100	(4.9)	1.5	584	(6.3)	4.7
A/PI**	90	(1.2)	6.4	47	(2.3)	3.2	137	(1.5)	4.7
AI/AN††	145	(2.0)	29.5	37	(1.8)	7.3	182	(2.0)	18.2
Hispanic§§	396	(5.5)	9.8	63	(3.1)	1.7	459	(5.0)	6.0
Other¶¶	16	(0.2)	NR¶	4	(0.2)	NR¶	20	(0.2)	—
Unknown¶¶¶	6	(0.1)	NR¶	1	—	NR¶	7	(0.1)	NR¶
Total	7,211	(100.0)	18.4	2,034	(100.0)	5.0	9,245	(100.0)	11.6
Marital Status***									
Married	2,721	(38.7)	NR†††	737	(37.1)	NR†††	3,458	(38.4)	NR†††
Never married	2,059	(29.3)	NR†††	431	(21.7)	NR†††	2,490	(27.6)	NR†††
Widowed	418	(6.0)	NR†††	176	(8.9)	NR†††	594	(6.6)	NR†††
Divorced	1,556	(22.1)	NR†††	580	(29.2)	NR†††	2,136	(23.7)	NR†††
Married, but separated	62	(0.9)	NR†††	17	(0.9)	NR†††	79	(0.9)	NR†††
Single, not otherwise specified	120	(1.7)	NR†††	24	(1.2)	NR†††	144	(1.6)	NR†††
Unknown	90	(1.3)	NR†††	23	(1.2)	NR†††	113	(1.3)	NR†††
Total	7,026	(100.0)	NR†††	1,988	(100.0)	NR†††	9,014	(100.0)	NR†††

* Percentages might not total 100% because of rounding.

† Per 100,000 population.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Rates not reported when number of decedents is <20.

** Asian/Pacific Islander.

†† American Indian/Alaskan Native.

§§ Includes persons of any race.

¶¶ Rates not computed for "other" or "unknown" categories.

*** Includes only decedents aged >18 years.

††† Rates cannot be computed for marital status because denominators are unknown.

TABLE 5. Number and percentage* of suicides, by sex of victim, method used, and location in which injury occurred — National Violent Death Reporting System, 16 states,† 2007

Characteristic	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Method						
Firearm	4,034	(56.0)	649	(31.9)	4,683	(50.7)
Sharp instrument	134	(1.9)	21	(1.0)	155	(1.7)
Blunt instrument	5	(0.1)	1	—	6	(0.1)
Poisoning	913	(12.7)	829	(40.8)	1,742	(18.8)
Hanging/Strangulation/Suffocation	1,756	(24.4)	376	(18.5)	2,132	(23.1)
Fall	107	(1.5)	41	(2.0)	148	(1.6)
Drowning	58	(0.8)	38	(1.9)	96	(1.0)
Fire/Burns	21	(0.3)	9	(0.4)	30	(0.3)
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	82	(1.1)	22	(1.1)	104	(1.1)
Other (single method)	14	(0.2)	6	(0.3)	20	(0.2)
Firearm and poisonings§	—	—	1	—	1	—
Firearm and other method type§	4	(0.1)	1	—	5	(0.1)
Poisoning and other method type§	12	(0.2)	16	(0.8)	28	(0.3)
Other combination of methods§	11	(0.2)	1	—	12	(0.1)
Unknown	60	(0.8)	23	(1.1)	83	(0.9)
Total	7,211	(100.0)	2,034	(100.0)	9,245	(100.0)
Location						
House	5,474	(75.9)	1,667	(82.0)	7,141	(77.2)
Street/Highway	236	(3.3)	43	(2.1)	279	(3.0)
Motor vehicle	174	(2.4)	32	(1.6)	206	(2.2)
Bar/Nightclub	2	—	—	—	2	—
Commercial/Retail area	45	(0.6)	2	(0.1)	47	(0.5)
Industrial or construction area	28	(0.4)	1	—	29	(0.3)
Office building	31	(0.4)	3	(0.1)	34	(0.4)
Parking lot/Public garage	86	(1.2)	30	(1.5)	116	(1.3)
Abandoned house, building or warehouse	4	(0.1)	—	—	4	—
Park, playground, sports/Athletic area	113	(1.6)	22	(1.1)	135	(1.5)
Preschool/School/College/School bus	15	(0.2)	6	(0.3)	21	(0.2)
Public transportation/Station/Railroad tracks	32	(0.4)	13	(0.6)	45	(0.5)
Hospital or medical facility	32	(0.4)	9	(0.4)	41	(0.4)
Supervised residential facility	40	(0.6)	7	(0.3)	47	(0.5)
Jail/Prison	115	(1.6)	5	(0.2)	120	(1.3)
Farm	30	(0.4)	1	—	31	(0.3)
Natural area	337	(4.7)	72	(3.5)	409	(4.4)
Hotel/Motel	115	(1.6)	38	(1.9)	153	(1.7)
Other	212	(2.9)	37	(1.8)	249	(2.7)
Unknown	90	(1.2)	46	(2.3)	136	(1.5)
Total	7,211	(100.0)	2,034	(100.0)	9,245	(100.0)

* Percentages might not total 100% because of rounding.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ Deaths involving more than one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 6. Number* and percentage of suicide victims tested for alcohol and drugs whose results were positive, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)§	6,366	(68.9)	2,194	(34.5)
BAC ≤0.08 g/dL§			723	(33.0)
BAC >0.08 g/dL§			1,361	(62.0)
Alcohol-positive, level unknown			110	(5.0)
Amphetamines	4,089	(44.2)	145	(3.6)
Antidepressants	3,678	(39.8)	1,101	(30.0)
Cocaine	4,361	(47.2)	386	(8.9)
Marijuana	3,229	(35.7)	246	(7.5)
Opiates	4,352	(47.1)	994	(22.8)
Other Drug(s)	3,905	(32.2)	2,045	(52.4)

* N = 9,245.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ BAC of 0.08% g/dL used as the standard for intoxication. Other substances indicated if any results were positive; levels for these substances are not measured.

TABLE 7. Number* and percentage† of suicides, by sex and associated circumstances — National Violent Death Reporting System, 16 states,§ 2007

Associated circumstances	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Mental health/Substance abuse						
Current depressed mood	2,547	(41.5)	755	(41.7)	3,302	(41.5)
Current mental health problem	2,433	(39.6)	1,139	(62.9)	3,572	(44.9)
Current mental health treatment	1,769	(28.8)	917	(50.6)	2,686	(33.8)
Alcohol problem	1,143	(18.6)	309	(17.1)	1,452	(18.3)
Other substance abuse problem	886	(14.4)	289	(16.0)	1,175	(14.8)
Interpersonal						
Intimate partner problem	1,948	(31.7)	479	(26.4)	2,427	(30.5)
Other relationship problem (nonintimate)	703	(11.4)	232	(12.8)	935	(11.8)
Suicide of family member or friend during preceding 5 yrs	114	(1.9)	33	(1.8)	147	(1.9)
Other death of family member or friend during preceding 5 yrs	387	(6.3)	122	(6.7)	509	(6.4)
Perpetrator of interpersonal violence during preceding month	345	(5.6)	29	(1.6)	374	(4.7)
Victim of interpersonal violence during preceding month	15	(0.2)	20	(1.1)	35	(0.4)
Life stressor						
Crisis during preceding 2 wks	2,005	(32.6)	483	(26.7)	2,488	(31.3)
Physical health problem	1,309	(21.3)	396	(21.9)	1,705	(21.4)
Job problem	774	(12.6)	141	(7.8)	915	(11.5)
Recent criminal legal problem	686	(11.2)	70	(3.9)	756	(9.5)
Noncriminal legal problem	238	(3.9)	56	(3.1)	294	(3.7)
Financial problem	747	(12.2)	182	(10.0)	929	(11.7)
School problem	69	(1.1)	15	(0.8)	84	(1.1)
Suicide event						
Left a suicide note	1,940	(31.6)	707	(39.0)	2,647	(33.3)
Disclosed intent to commit suicide	1,702	(27.7)	525	(29.0)	2,227	(28.0)
History of suicide attempt(s)	972	(15.8)	612	(33.8)	1,584	(19.9)

* N = 7,952 (6,141 males and 1,811 females). Circumstances unknown for 1,293 of 9,245 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 8. Number* and percent† of suicide decedents who had received a diagnosis of a current mental health problem, by diagnosis — National Violent Death Reporting System, 16 states,‡ 2007

Mental health problem	No.	(%)
Depression/Dysthymia	2,677	(74.9)
Bipolar disorder	518	(14.5)
Anxiety disorder	290	(8.1)
Schizophrenia	166	(4.6)
PTSD¶	60	(1.7)
OCD**	16	(0.4)
ADD/ADHD††	32	(0.9)
Eating disorder	7	(0.2)
Other	150	(4.2)
Unknown	308	(8.6)

* N = 3,572.

† Percentages might exceed 100% because multiple diagnosis categories might have been coded.

‡ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Posttraumatic stress disorder.

** Obsessive-compulsive disorder.

†† Attention deficit disorder/Attention deficit and hyperactivity disorder.

TABLE 9. Number,* percentage,† and rate‡ of homicides/legal intervention deaths, by method used and month in which death occurred — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	No.	(%)	Rate
Method			
Firearm	3,017	(66.1)	3.8
Sharp instrument	551	(12.1)	0.7
Blunt instrument	266	(5.8)	0.3
Poisoning	23	(0.5)	—
Hanging/Strangulation/Suffocation	138	(3.0)	0.2
Personal weapons (hands, feet, or fists)	155	(3.4)	0.2
Fall	17	(0.4)	NR**
Drowning	9	(0.2)	NR**
Fire/Burns	28	(0.6)	—
Shaking	28	(0.6)	—
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	35	(0.8)	—
Intentional neglect	15	(0.3)	NR**
Other (single method)	24	(0.5)	—
Firearm and other method type††	29	(0.6)	—
Poisoning and other method type††	7	(0.2)	NR**
Other combination of methods††	92	(2.0)	0.1
Unknown	129	(2.8)	0.2
Total	4,563	(100.0)	5.7
Month			
January	358	(7.8)	0.4
February	288	(6.3)	0.4
March	367	(8.0)	0.5
April	395	(8.7)	0.5
May	383	(8.4)	0.5
June	424	(9.3)	0.5
July	401	(8.8)	0.5
August	453	(9.9)	0.6
September	365	(8.0)	0.5
October	368	(8.1)	0.5
November	360	(7.9)	0.5
December	391	(8.6)	0.5
Unknown	10	(0.2)	NR**
Total	4,563	(100.0)	5.7

* No. victims = 4,545; no. suspects/victims = 18; no. live suspects = 4,136; total includes 4,545 victims and 18 suspects who were subsequently killed; no. incidents = 4,324.

† Percentages might not total 100% because of rounding.

‡ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rates not reported when number of decedents is <20.

†† Deaths involving more than one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 10. Number and percentage* of homicides/legal intervention deaths, by victim's marital status and relationship to suspect — National Violent Death Reporting System, 16 states,† 2007

Characteristic	No.	(%)
Marital Status§		
Married	919	(22.7)
Never married	2,144	(53.0)
Widowed	126	(3.1)
Divorced	528	(13.0)
Married, but separated	22	(0.5)
Single, not otherwise specified	222	(5.5)
Unknown	87	(2.1)
Total	4,048	(100.0)
Relationship		
Spouse/Intimate partner (current or former)	477	(10.5)
Parent	73	(1.6)
Child	153	(3.4)
Other intimate partner involvement¶	26	(0.6)
Other relative	120	(2.6)
Acquaintance/Friend	716	(15.7)
Rival gang member	35	(0.8)
Stranger	382	(8.4)
Victim injured by a law enforcement officer	153	(3.4)
Other specified relationship	335	(7.3)
More than one relationship mentioned	86	(1.9)
Multiple suspects in incident	127	(2.8)
Relationship unknown/missing	1,880	(41.2)
Total	4,563	(100.0)

* Percentages might not total 100% because of rounding.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ Includes only those victims aged >18 years.

¶ Death because of intimate partner-related violence, but not between the intimate partners themselves (e.g., when child is killed by the mother's partner or the teenager kills his mother's partner).

TABLE 11. Number, percentage,* and rate† of homicides/legal intervention deaths, by victim's sex, age group, and race/ethnicity — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	Male			Female			Total		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Age group (yrs)									
<1	64	(1.8)	11.1	45	(4.5)	8.2	109	(2.4)	9.7
1–4	51	(1.4)	2.3	45	(4.5)	2.1	96	(2.1)	2.2
5–9	21	(0.6)	0.8	25	(2.5)	1.0	46	(1.0)	0.9
10–14	30	(0.8)	1.1	24	(2.4)	0.9	54	(1.2)	1.0
15–19	453	(12.7)	15.8	77	(7.6)	2.8	530	(11.6)	9.5
20–24	699	(19.7)	24.5	113	(11.2)	4.2	812	(17.8)	14.7
25–29	584	(16.4)	20.5	108	(10.7)	3.9	692	(15.2)	12.4
30–34	392	(11.0)	14.9	74	(7.3)	2.9	466	(10.2)	8.9
35–44	542	(15.3)	9.3	198	(19.6)	3.4	740	(16.2)	6.3
45–54	405	(11.4)	7.0	161	(16.0)	2.7	566	(12.4)	4.8
55–64	191	(5.4)	4.5	65	(6.4)	1.4	256	(5.6)	2.9
65–74	72	(2.0)	3.1	34	(3.4)	1.2	106	(2.3)	2.1
75–84	40	(1.1)	3.1	28	(2.8)	1.4	68	(1.5)	2.1
≥85	8	(0.2)	NR¶	12	(1.2)	NR¶	20	(0.4)	1.5
Unknown	2	(0.1)	NR¶	—	—	NR¶	2	—	NR¶
Total	3,554	(100.0)	9.0	1,009	(100.0)	2.5	4,563	(100.0)	5.7
Race/ethnicity									
White, non-Hispanic	960	(27.0)	3.5	485	(48.1)	1.7	1,445	(31.7)	2.6
Black, non-Hispanic	2,004	(56.4)	34.3	368	(36.5)	5.7	2,372	(52.0)	19.3
A/PI**	40	(1.1)	2.8	23	(2.3)	1.5	63	(1.4)	2.2
AI/AN††	79	(2.2)	16.1	26	(2.6)	5.1	105	(2.3)	10.5
Hispanic§§	447	(12.6)	11.0	103	(10.2)	2.9	550	(12.1)	7.2
Other	21	(0.6)	0.1	3	(0.3)	NR¶	24	(0.5)	—
Unknown	3	(0.1)	NR¶	1	(0.1)	NR¶	4	(0.1)	NR¶
Total	3,554	(100.0)	9.0	1,009	(100.0)	2.5	4,563	(100.0)	5.7

* Percentages might not total 100% because of rounding.

† Per 100,000 population.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Rates not reported when number of decedents is <20.

** Asian/Pacific Islander.

†† American Indian/Alaskan Native.

§§ Includes persons of any race.

TABLE 12. Number and percentage* of homicides/legal intervention deaths, by victim's sex, method used, and location in which injury occurred — National Violent Death Reporting System, 16 states,† 2007

Method/Location	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Method						
Firearm	2,549	(71.7)	468	(46.4)	3,017	(66.1)
Sharp instrument	385	(10.8)	166	(16.5)	551	(12.1)
Blunt instrument	192	(5.4)	74	(7.3)	266	(5.8)
Poisoning	11	(0.3)	12	(1.2)	23	(0.5)
Hanging/Strangulation/Suffocation	51	(1.4)	87	(8.6)	138	(3.0)
Personal weapons (hands, feet, or fists)	108	(3.0)	47	(4.7)	155	(3.4)
Fall	14	(0.4)	3	(0.3)	17	(0.4)
Drowning	3	(0.1)	6	(0.6)	9	(0.2)
Fire/Burns	17	(0.5)	11	(1.1)	28	(0.6)
Shaking	18	(0.5)	10	(1.0)	28	(0.6)
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	26	(0.7)	9	(0.9)	35	(0.8)
Intentional neglect	8	(0.2)	7	(0.7)	15	(0.3)
Other (single method)	14	(0.4)	10	(1.0)	24	(0.5)
Firearm and other method type§	22	(0.6)	7	(0.7)	29	(0.6)
Poisoning and other method type§	2	(0.1)	5	(0.5)	7	(0.2)
Other combination of methods§	43	(1.2)	49	(4.9)	92	(2.0)
Unknown	91	(2.6)	38	(3.8)	129	(2.8)
Total	3,554	(100.0)	1,009	(100.0)	4,563	(100.0)
Location						
House	1,590	(44.7)	733	(72.6)	2,323	(50.9)
Street/Highway	923	(26.0)	69	(6.8)	992	(21.7)
Motor vehicle	141	(4.0)	19	(1.9)	160	(3.5)
Bar/Nightclub	65	(1.8)	1	(0.1)	66	(1.4)
Commercial/Retail area	107	(3.0)	27	(2.7)	134	(2.9)
Industrial or construction area	11	(0.3)	3	(0.3)	14	(0.3)
Office building	12	(0.3)	2	(0.2)	14	(0.3)
Parking lot/Public garage	192	(5.4)	19	(1.9)	211	(4.6)
Abandoned house, building, or warehouse	11	(0.3)	1	(0.1)	12	(0.3)
Park, playground, sports/athletic area	91	(2.6)	18	(1.8)	109	(2.4)
Preschool/School/College/School bus	19	(0.5)	14	(1.4)	33	(0.7)
Public transportation/Station/Railroad tracks	2	(0.1)	—	—	2	—
Hospital or medical facility	9	(0.3)	3	(0.3)	12	(0.3)
Supervised residential facility	9	(0.3)	1	(0.1)	10	(0.2)
Jail/Prison	26	(0.7)	—	—	26	(0.6)
Farm	10	(0.3)	1	(0.1)	11	(0.2)
Natural area	97	(2.7)	27	(2.7)	124	(2.7)
Hotel/Motel	34	(1.0)	19	(1.9)	53	(1.2)
Other	119	(3.3)	24	(2.4)	143	(3.1)
Unknown	86	(2.4)	28	(2.8)	114	(2.5)
Total	3,554	(100.0)	1,009	(100)	4,563	(100.0)

* Percentages might not total 100% because of rounding.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ Deaths involving more than one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 13. Number* and percentage of homicide/legal intervention victims tested for alcohol and drugs whose results were positive, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology Variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC) [§]	3,556	(77.9)	1,231	(34.6)
BAC ≤ 0.08 g/dL [§]			428	(34.8)
BAC >0.08 g/dL [§]			682	(55.4)
Alcohol positive, level unknown			121	(9.8)
Amphetamines	2,481	(54.4)	157	(6.3)
Antidepressants	1,586	(34.8)	68	(4.3)
Cocaine	2,685	(58.8)	422	(15.7)
Marijuana	1,540	(33.8)	308	(20.0)
Opiates	2,392	(52.4)	162	(6.8)
Other drug(s)	2,024	(44.4)	413	(20.4)

* N = 4,563.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 14. Number* and percentage† of homicide/legal intervention deaths, by associated circumstances and victim's sex — National Violent Death Reporting System, 16 states,§ 2007

Circumstance	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Precipitated by another crime	790	(35.7)	172	(23.0)	962	(32.5)
Crime in progress [¶]	605	(76.6)	126	(73.3)	731	(76.0)
Argument about money/property	141	(6.4)	25	(3.3)	161	(5.4)
Jealousy (lovers' triangle)	79	(3.6)	40	(5.4)	119	(4.0)
Other argument, abuse, conflict	927	(41.9)	183	(24.5)	1,110	(37.5)
Drug involvement	333	(15.0)	61	(8.2)	394	(13.3)
Justifiable self defense/Law enforcement	230	(10.4)	6	(0.8)	236	(8.0)
Brawl	48	(2.2)	5	(0.7)	53	(1.8)
Mercy killing	1	—	3	(0.4)	4	(0.1)
Victim was a bystander	37	(1.7)	22	(2.9)	59	(2.0)
Victim was a police officer on duty	11	(0.5)	—	—	11	(0.4)
Victim was an intervener assisting a crime victim	17	(0.8)	5	(0.7)	22	(0.7)
Victim used a weapon	304	(13.7)	14	(1.9)	318	(10.7)
Intimate partner-violence-related	211	(9.5)	392	(52.5)	603	(20.4)
Hate crime	5	(0.2)	—	—	5	(0.2)
Mentally ill suspect	42	(1.9)	31	(4.1)	73	(2.5)
Drive-by shooting	95	(4.3)	10	(1.3)	105	(3.5)
Random violence	19	(0.9)	7	(0.9)	26	(0.9)
Gang-related	136	(6.1)	10	(1.3)	146	(4.9)

* N = 2,961 (2,214 males and 747 females). Circumstances were unknown for 1,602 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Denominator is only cases that were precipitated by another crime.

TABLE 15. Number and percentage* of homicides/legal intervention deaths precipitated by another crime in progress at the time of injury, by type of crime — National Violent Death Reporting System, 16 states,† 2007

Crime type	No.	(%)
Robbery	360	(37.4)
Burglary	89	(9.3)
Assault/Homicide	224	(23.3)
Rape, sexual assault	32	(3.3)
Motor vehicle theft	29	(3.0)
Arson	11	(1.1)
Drug trade	79	(8.2)
Witness intimidation/Elimination	3	(0.3)
Gambling	4	(0.4)
Other	58	(6.0)
Unknown	54	(5.6)

* Percentages might exceed 100% because multiple crimes might have been coded.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 16. Number,* percentage,† and rate§ of undetermined deaths¶ by method used, and month in which death occurred — National Violent Death Reporting System, 16 states, 2007**

Characteristic	No.	(%)	Rate
Method			
Firearm	64	(2.7)	0.1
Sharp instrument	6	(0.2)	NR††
Blunt instrument	49	(2.0)	0.1
Poisoning	1,563	(65.0)	2.0
Hanging/Strangulation/Suffocation	42	(1.7)	0.1
Personal weapons (hands, feet, fists)	6	(0.2)	NR††
Fall	40	(1.7)	0.1
Drowning	51	(2.1)	0.1
Fire/Burns	19	(0.8)	NR††
Shaking	—	—	NR††
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	30	(1.2)	—
Intentional neglect	4	(0.2)	NR††
Other (single method)	95	(4.0)	0.1
Firearm and other method type§§	1	(0.0)	NR††
Poisoning and other method type§§	21	(0.9)	—
Other combination of methods§§	3	(0.1)	NR††
Unknown	409	(17.0)	0.5
Total	2,403	(100.0)	3.0
Month			
January	195	(8.1)	0.2
February	208	(8.7)	0.3
March	203	(8.4)	0.3
April	193	(8.0)	0.2
May	201	(8.4)	0.3
June	208	(8.7)	0.3
July	214	(8.9)	0.3
August	217	(9.0)	0.3
September	185	(7.7)	0.2
October	184	(7.7)	0.2
November	185	(7.7)	0.2
December	206	(8.6)	0.3
Unknown	4	(0.2)	NR††
Total	2,403	(100.0)	3.0

* No. victims = 2,403; no. suspects/victims = 0; no. live suspects = 24; no. incidents = 2,389 plus three incidents categorized in "other combinations of deaths" in Table 1.

† Percentages might not total 100% because of rounding.

§ Per 100,000 population.

¶ Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

** Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

†† Rates not reported when number of decedents is <20.

§§ Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 17. Number, percentage* and rate† of undetermined deaths,§ by victim's sex, age group, race/ethnicity, and marital status — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	Male			Female			Total		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Age group (yrs)									
<1	152	(10.2)	26.4	96	(10.5)	17.5	248	(10.3)	22.0
1–4	16	(1.1)	NR**	14	(1.5)	NR**	30	(1.2)	0.7
5–9	7	(0.5)	NR**	5	(0.5)	NR**	12	(0.5)	NR**
10–14	8	(0.5)	NR**	4	(0.4)	NR**	12	(0.5)	NR**
15–19	59	(4.0)	2.1	12	(1.3)	NR**	71	(3.0)	1.3
20–24	106	(7.1)	3.7	47	(5.1)	1.8	153	(6.4)	2.8
25–29	116	(7.8)	4.1	58	(6.3)	2.1	174	(7.2)	3.1
30–34	113	(7.6)	4.3	67	(7.3)	2.6	180	(7.5)	3.4
35–44	339	(22.8)	5.8	198	(21.7)	3.4	537	(22.3)	4.6
45–54	365	(24.5)	6.3	279	(30.5)	4.6	644	(26.8)	5.5
55–64	146	(9.8)	3.4	88	(9.6)	1.9	234	(9.7)	2.7
65–74	28	(1.9)	1.2	25	(2.7)	0.9	53	(2.2)	1.1
75–84	20	(1.3)	1.5	12	(1.3)	NR**	32	(1.3)	1.0
≥85	6	(0.4)	NR**	9	(1.0)	NR**	15	(0.6)	NR**
Unknown	6	(0.4)	NR**	2	(0.2)	NR**	8	(0.3)	NR**
Total	1,487	(100.0)	3.8	916	(100.0)	2.3	2,403	(100.0)	3.0
Race/ethnicity									
White, non-Hispanic	1,027	(69.1)	3.7	692	(75.5)	2.4	1,719	(71.5)	3.1
Black, non-Hispanic	312	(21.0)	5.3	149	(16.3)	2.3	461	(19.2)	3.7
A/PI††	8	(0.5)	NR**	5	(0.5)	NR**	13	(0.5)	NR**
AI/AN§§	37	(2.5)	7.5	24	(2.6)	4.7	61	(2.5)	6.1
Hispanic¶¶	95	(6.4)	2.3	44	(4.8)	1.2	139	(5.8)	1.8
Other	8	(0.5)	NR**	2	(0.2)	NR**	10	(0.4)	NR**
Total	1,487	(100.0)	3.8	916	(100.0)	2.3	2,403	(100.0)	3.0
Marital Status***									
Married	307	(24.1)	NR†††	270	(34.1)	NR†††	577	(27.9)	NR†††
Never married	587	(46.1)	NR†††	225	(28.4)	NR†††	812	(39.3)	NR†††
Widowed	33	(2.6)	NR†††	54	(6.8)	NR†††	87	(4.2)	NR†††
Divorced	284	(22.3)	NR†††	219	(27.7)	NR†††	503	(24.3)	NR†††
Married, but separated	4	(0.3)	NR†††	5	(0.6)	NR†††	9	(0.4)	NR†††
Single, not otherwise specified	10	(0.8)	NR†††	5	(0.6)	NR†††	15	(0.7)	NR†††
Unknown	49	(3.8)	NR†††	13	(1.6)	NR†††	65	(3.1)	NR†††
Total	1,274	(100.0)	NR†††	791	(100.0)	NR†††	2,068	(100.0)	NR†††

* Percentages might not total 100% because of rounding.

† Per 100,000 population.

§ Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rate not reported when number of decedents is <20.

†† Asian/Pacific Islander.

§§ American Indian/Alaskan Native.

¶¶ Includes persons of all races.

*** Includes only those decedents aged >18 years.

††† Rates cannot be computed for marital status because denominators are unknown.

TABLE 18. Number and percentage* of undetermined deaths,† by decedent's sex, method used, and location in which injury occurred — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Method						
Firearm	48	(3.2)	16	(1.7)	64	(2.7)
Sharp instrument	3	(0.2)	3	(0.3)	6	(0.2)
Blunt instrument	33	(2.2)	16	(1.7)	49	(2.0)
Poisoning	946	(63.6)	617	(67.4)	1,563	(65.0)
Hanging/Strangulation/Suffocation	30	(2.0)	12	(1.3)	42	(1.7)
Personal weapons (hands, feet, fists)	4	(0.3)	2	(0.2)	6	(0.2)
Fall	21	(1.4)	19	(2.1)	40	(1.7)
Drowning	33	(2.2)	18	(2.0)	51	(2.1)
Fire/Burns	13	(0.9)	6	(0.7)	19	(0.8)
Shaking	—	—	8	(0.9)	8	(0.3)
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	22	(1.5)	—	—	22	(0.9)
Intentional neglect	—	—	4	(0.4)	4	(0.2)
Other (single method)	62	(4.2)	33	(3.6)	95	(4.0)
Firearm and other method type¶	1	(0.1)	—	—	1	—
Poisoning and other method type¶	14	(0.9)	7	(0.8)	21	(0.9)
Other combination of methods¶	1	(0.1)	2	(0.2)	3	(0.1)
Unknown	256	(17.2)	153	(16.7)	409	(17.0)
Total	1,487	(100.0)	916	(100.0)	2,403	(100.0)
Location						
House	1,116	(75.1)	757	(82.6)	1,873	(77.9)
Street/Highway	50	(3.4)	21	(2.3)	71	(3.0)
Motor vehicle	16	(1.1)	7	(0.8)	23	(1.0)
Bar/Nightclub	—	—	1	(0.1)	1	—
Commercial/Retail area	5	(0.3)	2	(0.2)	7	(0.3)
Industrial or construction area	5	(0.3)	1	(0.1)	6	(0.2)
Office building	1	(0.1)	—	—	1	—
Parking lot/Public garage	5	(0.3)	5	(0.5)	10	(0.4)
Abandoned house, building or warehouse	5	(0.3)	2	(0.2)	7	(0.3)
Park, playground, sports/Athletic area	7	(0.5)	4	(0.4)	11	(0.5)
Preschool/School/College/School bus	2	(0.1)	—	—	2	(0.1)
Public transportation/Station/Railroad tracks	8	(0.5)	1	(0.1)	9	(0.4)
Hospital or medical facility	8	(0.5)	7	(0.8)	15	(0.6)
Supervised residential facility	28	(1.9)	8	(0.9)	36	(1.5)
Jail/Prison	6	(0.4)	1	(0.1)	7	(0.3)
Farm	4	(0.3)	1	(0.1)	5	(0.2)
Natural area	53	(3.6)	23	(2.5)	76	(3.2)
Hotel/Motel	22	(1.5)	6	(0.7)	28	(1.2)
Other	68	(4.6)	15	(1.6)	83	(3.5)
Unknown	78	(5.2)	54	(5.9)	132	(5.5)
Total	1,487	(100.0)	916	(100.0)	2,403	(100.0)

* Percentages might not total 100% because of rounding.

† Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 19. Number* and percentage of victims of undetermined intent† tested for alcohol and drugs whose results were positive, by toxicology variable — National Violent Death Reporting System, 16 states,§ 2007

Toxicology variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)¶	2,051	(85.4)	578	(28.2)
BAC ≤0.08 g/dL¶			227	(39.3)
BAC >0.08 g/dL¶			319	(55.2)
Alcohol positive, level unknown			32	(5.5)
Amphetamines	1,941	(80.8)	89	(4.6)
Antidepressants	1,740	(72.4)	484	(27.8)
Cocaine	2,005	(83.4)	423	(21.1)
Marijuana	1,077	(44.8)	101	(9.4)
Opiates	2,040	(84.9)	1,154	(56.6)
Other drug(s)	1,948	(81.1)	1,059	(54.4)

* N = 2,403.

† Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 20. Number* and percentage† of deaths of undetermined intent,§ by victim's sex and associated circumstances — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Mental health/Substance abuse						
Current depressed mood	117	(10.8)	111	(16.4)	228	(13.0)
Current mental health problem	335	(30.9)	351	(52.0)	686	(39.0)
Current mental health treatment	257	(23.7)	301	(44.6)	558	(31.7)
Alcohol problem	359	(33.1)	146	(21.6)	505	(28.7)
Other substance abuse problem	706	(65.1)	391	(57.9)	1,097	(62.4)
Interpersonal						
Intimate partner problem	97	(8.9)	93	(13.8)	190	(10.8)
Other relationship problem (nonintimate)	49	(4.5)	48	(7.1)	97	(5.5)
Suicide of family member or friend within past 5 years	3	(0.3)	4	(0.6)	7	(0.4)
Other death of family member or friend within past 5 years	36	(3.3)	46	(6.8)	82	(4.7)
Perpetrator of interpersonal violence within past month	16	(1.5)	3	(0.4)	19	(1.1)
Victim of interpersonal violence within past month	9	(0.8)	14	(2.1)	23	(1.3)
Life stressor						
Crisis in past two weeks	148	(13.7)	115	(17.0)	263	(15.0)
Physical health problem	275	(25.4)	235	(34.8)	510	(29.0)
Job problem	46	(4.2)	17	(2.5)	63	(3.6)
Recent criminal legal problem	47	(4.3)	17	(2.5)	64	(3.6)
Non-criminal legal problem	21	(1.9)	10	(1.5)	31	(1.8)
Financial problem	27	(2.5)	22	(3.3)	49	(2.8)
School problem	3	(0.3)	1	(0.1)	4	(0.2)
Suicide event						
Left a suicide note	12	(1.1)	17	(2.5)	29	(1.6)
Disclosed intent to commit suicide	75	(6.9)	51	(7.6)	126	(7.2)
History of suicide attempt(s)	91	(8.4)	93	(13.8)	184	(10.5)

* N = 1,759 (1,084 males and 675 females). Circumstances were unknown for 644 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 21. Number* and percentage† of victims of undetermined intent§ who had received a diagnosis of a current mental health problem, by diagnosis — National Violent Death Reporting System, 16 states,¶ 2007

Mental health problem	No.	(%)
Depression/Dysthymia	420	(61.2)
Bipolar disorder	110	(16.0)
Anxiety disorder	86	(12.5)
Schizophrenia	40	(5.8)
PTSD**	14	(2.0)
OCD††	4	(0.6)
ADD/ADHD§§	8	(1.2)
Eating disorder	—	—
Other	30	(4.4)
Unknown	81	(11.8)

* N = 686.

† Percentages might exceed 100% because two or more diagnosis categories per person could be coded.

§ Deaths that result from the use of force or power against oneself or another person for which evidence indicating one manner of death is no more compelling than evidence indicating another.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Posttraumatic stress disorder.

†† Obsessive compulsive disorder.

§§ Attention deficit disorder/hyperactivity disorder.

TABLE 22. Number* and percentage† of unintentional firearm deaths, by victim's sex, race/ethnicity, age group, month in which the death occurred, and location of injury — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	No.	(%)
Sex		
Male	97	(90.7)
Female	10	(9.3)
Total	107	(100)
Race/ethnicity		
White, non-Hispanic	73	(68.2)
Black, non-Hispanic	24	(22.4)
A/PI¶	—	—
AI/AN**	3	(2.8)
Hispanic††	7	(6.5)
Total	107	(100.0)
Age		
<1	—	—
1–4	4	(3.7)
5–9	6	(5.6)
10–14	6	(5.6)
15–19	21	(19.6)
20–24	12	(11.2)
25–29	9	(8.4)
30–34	3	(2.8)

TABLE 22. (Continued) Number* and percentage† of unintentional firearm deaths, by victim's sex, race/ethnicity, age group, month in which the death occurred, and location of injury — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	No.	(%)
Age		
35–44	11	(10.3)
45–54	9	(8.4)
55–64	11	(10.3)
65–74	6	(5.6)
75–84	5	(4.7)
≥85	4	(3.7)
Total	107	(100.0)
Month		
January	10	(9.3)
February	10	(9.3)
March	4	(3.7)
April	9	(8.4)
May	8	(7.5)
June	4	(3.7)
July	9	(8.4)
August	9	(8.4)
September	10	(9.3)
October	9	(8.4)
November	19	(17.8)
December	5	(4.7)
Unknown	1	(0.9)
Total	107	(100.0)
Location		
House	69	(64.5)
Street/Highway	2	(1.9)
Motor vehicle	3	(2.8)
Industrial/Construction area	1	(0.9)
Supervised residential facility	1	(0.9)
Farm	1	(0.9)
Natural area	18	(16.8)
Other§§	4	(3.7)
Unknown	8	(7.5)
Total	107	(100.0)
Firearm Type		
Handgun	44	(41.1)
Shotgun	19	(17.8)
Rifle	21	(19.6)
Other firearm	2	(1.9)
Unknown	21	(19.6)
Total	107	(100.0)

* No. incidents = 107; no. decedents = 107; no. live suspects = 35.

† Percentages might not total 100% because of rounding.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Asian/Pacific Islander.

** American Indian/Alaskan Native.

†† Includes persons of any race.

§§ Includes military training exercise, private land campsites, and private hunting land attached to homes.

TABLE 23. Number* and percentage† of unintentional firearm deaths, by context and circumstances of injury — National Violent Death Reporting System, 16 states,§ 2007

Circumstances	No.	(%)
Context of Injury		
Hunting	19	(24.7)
Target shooting	3	(3.9)
Loading/Unloading gun	8	(10.4)
Cleaning gun	5	(6.5)
Showing gun to others	11	(14.3)
Playing with gun	23	(29.9)
Other context of injury	16	(20.8)
Circumstances of injury		
Thought safety was engaged	1	(1.3)
Thought gun was unloaded, magazine disengaged	10	(13.0)
Thought gun was unloaded, other	10	(13.0)
Unintentionally pulled trigger	15	(19.5)
Bullet ricochet	1	(1.3)
Gun defect or malfunction	4	(5.2)
Dropped gun	1	(1.3)
Gun mistaken for toy	1	(1.3)
Other mechanism of injury	20	(26.0)

* N = 77. Circumstances were unknown for 30 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 24. Number* and percentage† of incidents involving multiple violent deaths, by incident type and method used — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	No.	(%)
Incident type		
Multiple suicides	12	(3.5)
Multiple homicides	148	(42.7)
Homicide followed by suicide	172	(49.6)
Other combinations of deaths	6	(1.7)
Undetermined	9	(2.6)
Total	347	(100.0)
Method		
Firearm	578	(73.7)
Sharp instrument	41	(5.2)
Blunt instrument	26	(3.3)
Poisoning	45	(5.7)
Hanging/Strangulation/Suffocation	41	(5.2)
Personal weapons	3	(0.4)
Drowning	6	(0.8)
Fire/Burns	21	(2.7)
Motor vehicles (e.g., buses, motorcycles, and other transport vehicles)	1	(0.1)
Other, single weapon	5	(0.6)
Firearm and other method type¶	1	(0.1)
Poisoning and other method type¶	2	(0.3)
Other combination of methods¶	5	(0.6)
Unknown	9	(1.1)
Total	784	(100.0)

* No. victims = 784; no. suspects = 368; no. incidents = 347; number of decedents includes 191 homicide suspects who subsequently killed themselves.

† Percentages might not total 100% because of rounding.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 25. Number, percentage,* and rate† of violent death incidents involving multiple victims, by victim's sex, race/ethnicity, and age group — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	Victims			Suspects¶	
	No.	(%)	Rate	No.	(%)
Sex					
Male	461	(58.8)	1.2	334	(90.8)
Female	323	(41.2)	0.8	23	(6.3)
Unknown	—	—	NR**	11	(3.0)
Total	784	(100.0)	0.9	368	(100.0)
Race/Ethnicity					
White, non-Hispanic	447	(57.0)	0.8	162	(44.0)
Black, non-Hispanic	208	(26.5)	1.7	108	(29.4)
A/PI††	25	(3.2)	0.9	8	(2.2)
AI/AN§§	19	(2.4)	NR**	9	(2.5)
Hispanic¶¶	82	(10.5)	1.1	37	(10.1)
Other	3	(0.4)	NR**	43	(11.7)
Unknown	—	—	NR**	1	(0.3)
Total	784	(100.0)	0.9	368	(100.0)
Age					
<1	5	(0.6)	NR**	—	—
1–4	22	(2.8)	0.5	—	—
5–9	23	(2.9)	0.4	—	—
10–14	19	(2.4)	NR**	—	—
15–19	72	(9.2)	1.3	28	(7.6)
20–24	95	(12.1)	1.7	43	(11.7)
25–29	95	(12.1)	1.7	40	(10.9)
30–34	68	(8.7)	1.3	36	(9.8)
35–44	147	(18.8)	1.3	70	(19.0)
45–54	113	(14.4)	1.0	48	(13.0)
55–64	52	(6.6)	0.6	25	(6.8)
65–74	29	(3.7)	0.6	6	(1.6)
75–84	33	(4.2)	1.0	10	(2.7)
≥85	9	(1.1)	NR**	2	(0.5)
Unknown	2	(0.3)	NR**	60	(16.3)
Total	784	(100.0)	0.9	368	(100.0)

* Percentages might not total 100% because of rounding.

† Per 100,000 population.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Number of suspects includes 166 homicide suspects who subsequently committed suicide. Rates cannot be computed for suspects because the number of suspects involved in an incident is not always known.

** Rates not reported when number of decedents is <20.

†† Asian/Pacific Islander.

§§ American Indian/Alaskan Native.

¶¶ Includes persons of any race.

TABLE 26. Number,* percentage,† and rate‡ of deaths involving a homicide followed by a suicide, by victim's sex, race/ethnicity, age group, and marital status — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	Homicide			Suicide		
	No.	(%)	Rate	No.	(%)	Rate
Sex						
Male	66	(27.5)	0.2	160	(93.0)	0.4
Female	174	(72.5)	0.4	12	(7.0)	NR**
Total	240	(100.0)	0.3	172	(100.0)	0.2
Race/ethnicity						
White, non-Hispanic	169	(70.4)	0.3	115	(66.9)	0.2
Black, non-Hispanic	33	(13.8)	0.3	27	(15.7)	0.2
A/PI††	9	(3.8)	NR**	7	(4.1)	NR**
AI/AN§§	4	(1.7)	NR**	3	(1.7)	NR**
Hispanic¶¶	22	(9.2)	0.3	20	(11.6)	0.3
Other	3	(1.3)	NR**	—	—	NR**
Total	240	(100.0)	0.3	172	(100.0)	0.2
Age group (yrs)						
<1	4	(1.7)	NR**	—	—	NR**
1–4	16	(6.7)	NR**	—	—	NR**
5–9	18	(7.5)	NR**	—	—	NR**
10–14	6	(2.5)	NR**	—	—	NR**
15–19	20	(8.3)	0.4	4	(2.3)	NR**
20–24	23	(9.6)	0.4	12	(7.0)	NR**
25–29	28	(11.7)	0.5	13	(7.6)	NR**
30–34	15	(6.3)	NR**	20	(11.6)	0.4
35–44	46	(19.2)	0.4	54	(31.4)	0.5
45–54	30	(12.5)	0.3	31	(18.0)	0.3
55–64	10	(4.2)	NR**	20	(11.6)	0.2
65–74	9	(3.8)	NR**	6	(3.5)	NR**
75–84	12	(5.0)	NR**	10	(5.8)	NR**
≥85	3	(1.3)	NR**	2	(1.2)	NR**
Unknown	—	—	NR**	—	—	NR**
Total	240	(100.0)	0.3	172	(100.0)	0.2
Marital Status***						
Married	66	(34.7)	NR†††	55	(32.4)	NR†††
Never married	60	(31.6)	NR†††	44	(25.9)	NR†††
Widowed	22	(11.6)	NR†††	36	(21.2)	NR†††
Divorced	34	(17.9)	NR†††	27	(15.9)	NR†††
Married, but separated	4	(2.1)	NR†††	3	(1.8)	NR†††
Single, not otherwise specified	3	(1.6)	NR†††	2	(1.2)	NR†††
Unknown	1	(0.5)	NR†††	3	(1.8)	NR†††
Total	190	(100.0)	NR†††	170	(100.0)	NR†††

* N = 172.

† Percentages might not total 100% because of rounding.

‡ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rates not reported when number of decedents is <20.

†† Asian/Pacific Islander.

§§ American Indian/Alaskan Native.

¶¶ Includes persons of any race.

*** Includes only decedents age >18 years.

††† Rates for marital status cannot be computed because denominators are unknown.

TABLE 27. Number and percentage* of homicides followed by suicide, by location in which injury occurred and method used — National Violent Death Reporting System, 16 states,† 2007

Characteristic	Decedent			
	Homicide		Suicide	
	No.	(%)	No.	(%)
Location				
House	181	(75.4)	136	(79.1)
Street/Highway	5	(2.1)	7	(4.1)
Motor vehicle	4	(1.7)	2	(1.2)
Commercial/Retail area	5	(2.1)	2	(1.2)
Industrial or construction area	—	—	1	(0.6)
Office building	2	(0.8)	4	(2.3)
Parking lot/public garage	4	(1.7)	4	(2.3)
Park, playground, sports/Athletic area	4	(1.7)	—	—
Preschool/School/College/School bus	24	(10.0)	1	(0.6)
Public transportation/Station/Railroad tracks	—	—	1	(0.6)
Hospital or medical facility	1	(0.4)	1	(0.6)
Supervised residential facility	—	—	1	(0.6)
Natural area	3	(1.3)	4	(2.3)
Hotel/Motel	1	(0.4)	2	(1.2)
Other	6	(2.5)	5	(2.9)
Unknown	—	—	1	(0.6)
Total	240	(100.0)	172	(100.0)
Method				
Firearm	192	(80.0)	138	(80.2)
Sharp instrument	6	(2.5)	5	(2.9)
Blunt instrument	6	(2.5)	1	(0.6)
Poisoning	5	(2.1)	8	(4.7)
Hanging/Strangulation/Suffocation	17	(7.1)	11	(6.4)
Personal weapons	2	(0.8)	—	—
Drowning	1	(0.4)	1	(0.6)
Fire/Burns	7	(2.9)	3	(1.7)
Motor vehicle	—	—	1	(0.6)
Other, single weapon	2	(0.8)	1	(0.6)
Poisoning and other method type§	—	—	1	(0.6)
Other combination of methods§	—	—	2	(1.2)
Unknown	2	(0.8)	—	—
Total	240	(100.0)	172	(100.0)

* Percentages might not total 100% because of rounding.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 28. Number* and percentage of homicides followed by suicide, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology variable	Homicide Victim				Suicide Victim			
	Tested		Positive		Tested		Positive	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)§	191	(79.6)	35	(18.3)	125	(72.7)	41	(32.8)
BAC ≤ 0.08 g/dL§			15	(42.9)			17	(41.5)
BAC >0.08 g/dL§			17	(48.6)			22	(53.7)
Alcohol positive, level unknown			3	(8.6)			2	(4.9)
Amphetamines	109	(45.4)	6	(5.5)	77	(44.8)	—	—
Antidepressants	86	(35.8)	6	(7.0)	55	(32.0)	6	(10.9)
Cocaine	115	(47.9)	3	(2.6)	84	(48.8)	6	(7.1)
Marijuana	77	(32.1)	2	(2.6)	54	(31.4)	3	(5.6)
Opiates	106	(44.2)	10	(9.4)	75	(43.6)	9	(12.0)
Other Drug(s)	95	(39.6)	30	(31.6)	64	(37.2)	21	(32.8)

* N = 240 homicide victims and 172 suicide victims.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 29. Number* and percentage† of homicide suspects who killed themselves after committing a homicide, by suicide circumstances — National Violent Death Reporting System, 16 states,§ 2007

Circumstance	No.	(%)
Mental health/Substance abuse		
Current depressed mood	14	(8.3)
Current mental health problem	11	(6.6)
Current mental health treatment	6	(3.6)
Alcohol problem	7	(4.2)
Other substance abuse problem	14	(8.3)
Interpersonal		
Intimate partner problem	136	(81.0)
Other relationship problem (nonintimate)	23	(13.7)
Suicide of family member or friend within past 5 years	1	(0.6)
Other death of family member or friend within past 5 years	21	(12.5)
Perpetrator of interpersonal violence within past month	128	(76.2)
Victim of interpersonal violence within past month	—	—
Life stressor		
Crisis in past two weeks	153	(91.1)
Physical health problem	11	(6.6)
Job problem	4	(2.4)
Recent criminal legal problem	32	(19.1)
Non-criminal legal problem	5	(3.0)
Financial problem	7	(4.2)
School problem	1	(0.6)
Suicide event		
Left a suicide note	29	(17.3)
Disclosed intent to commit suicide	10	(6.0)
History of suicide attempt(s)	3	(1.8)

* N = 168. Circumstances were unknown for four deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 30. Number,* percentage,† and rate‡ of deaths related to violence by intimate partners, by victim's and suspect's sex, race/ethnicity, age group, and marital status — National Violent Death Reporting System, 16 states,¶ 2007

Characteristic	Victim			Suspect**	
	No	(%)	Rate	No	(%)
Sex					
Male*	218	(35.6)	0.6	451	(77.8)
Female	394	(64.4)	1.0	122	(21.0)
Unknown	—	—	NR††	7	(1.2)
Total	612	(100.0)	0.8	580	(100.0)
Race/ethnicity					
White, non-Hispanic	314	(51.3)	0.6	252	(43.5)
Black, non-Hispanic	200	(32.7)	1.6	186	(32.1)
A/PI§§	16	(2.6)	NR††	11	(1.9)
AI/AN¶¶	21	(3.4)	2.1	15	(2.6)
Hispanic***	60	(9.8)	0.8	58	(10.0)
Other	1	(0.2)	NR††	53	(9.1)
Unknown	—	—	NR††	5	(0.9)
Total	612	(100.0)	0.8	580	(100.0)
Age group (yrs)					
<1	6	(1.0)	NR††	—	—
1–4	12	(2.0)	NR††	—	—
5–9	9	(1.5)	NR††	—	—
10–14	5	(0.8)	NR††	—	—
15–19	38	(6.2)	0.7	23	(4.0)
20–24	72	(11.8)	1.3	52	(9.0)
25–29	67	(10.9)	1.2	60	(10.3)
30–34	61	(10.0)	1.2	57	(9.8)
35–44	160	(26.1)	1.4	136	(23.5)
45–54	118	(19.3)	1.0	97	(16.7)
55–64	39	(6.4)	0.4	47	(8.1)
65–74	16	(2.6)	NR††	16	(2.8)
75–84	5	(0.8)	NR††	7	(1.2)
≥85	4	(0.7)	NR††	3	(0.5)
Unknown	—	—	NR††	82	(14.1)
Total	612	(100.0)	0.8	580	(100.0)
Marital Status†††					
Married	213	(37.8)	NR §§§	NR ¶¶¶	NR ¶¶¶
Never Married	179	(31.8)	NR §§§	NR ¶¶¶	NR ¶¶¶
Widowed	30	(5.3)	NR §§§	NR ¶¶¶	NR ¶¶¶
Divorced	116	(20.6)	NR §§§	NR ¶¶¶	NR ¶¶¶
Married, but separated	8	(1.4)	NR §§§	NR ¶¶¶	NR ¶¶¶
Single, not otherwise specified	11	(2.0)	NR §§§	NR ¶¶¶	NR ¶¶¶
Unknown	6	(1.1)	NR §§§	NR ¶¶¶	NR ¶¶¶
Total	563	(100.0)	NR §§§	NR ¶¶¶	NR ¶¶¶

* N = 562. Number of male victims is nine more than Table 14 because their manner of death was undetermined.

† Percentages might not total 100% because of rounding.

‡ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rates cannot be calculated for suspects because the number of suspects involved in an incident is not always known.

†† Rates not reported when number of decedents is <20.

§§ Asian/Pacific Islander.

¶¶ American Indian/Alaska Native.

*** Includes persons of any race.

††† Includes only those decedents aged >18 years.

§§§ Rates for marital status cannot be computed because denominators are unknown.

¶¶¶ Data not available.

TABLE 31. Number* and percentage of deaths by intimate partner violence, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)§	499	(81.5)	169	(33.9)
BAC ≤0.08 g/dL§			55	(32.5)
BAC >0.08 g/dL§			102	(60.4)
Alcohol positive, level unknown			12	(7.1)
Amphetamines	320	(52.3)	11	(3.4)
Antidepressants	231	(37.8)	17	(7.4)
Cocaine	346	(56.5)	44	(12.7)
Marijuana	231	(37.8)	18	(7.8)
Opiates	318	(52.0)	19	(6.0)
Other Drug(s)	264	(43.1)	62	(23.5)

* N = 612.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 32. Number and percentage* of suicides among former or current military personnel, by sex, race/ethnicity, age group, marital status, and method used — National Violent Death Reporting System, 16 states,† 2007

Characteristic	No.	(%)
Sex of victim		
Male	1,713	(96.6)
Female	61	(3.4)
Total	1,774	(100.0)
Race/ethnicity		
White, non-Hispanic	1,627	(91.7)
Black, non-Hispanic	91	(5.1)
A/PI§	5	(0.3)
AI/AN¶	10	(0.6)
Hispanic**	41	(2.3)
Total	1,774	(100.0)
Age of victim		
<19	8	(0.5)
20–24	59	(3.3)
25–29	64	(3.6)
30–34	53	(3.0)
35–44	230	(13.0)
45–54	324	(18.3)
55–64	358	(20.2)
65–74	253	(14.3)
75–84	312	(17.6)
≥85	113	(6.4)
Total	1,774	(100.0)
Marital status††		
Married	832	(46.9)
Never married	242	(13.6)
Widowed	215	(12.1)

TABLE 32. (Continued) Number and percentage* of suicides among former or current military personnel, by sex, race/ethnicity, age group, marital status, and method used — National Violent Death Reporting System, 16 states,† 2007

Characteristic	No.	(%)
Marital status††		
Divorced	457	(25.8)
Married, but separated	8	(0.5)
Single, not otherwise specified	12	(0.7)
Unknown	7	(0.4)
Total	1,773	(100.0)
Method		
Firearm	1,193	(67.2)
Sharp instrument	27	(1.5)
Poisoning	218	(12.3)
Hanging/Strangulation/Suffocation	258	(14.5)
Fall	18	(1.0)
Drowning	8	(0.5)
Fire/Burns	4	(0.2)
Motor vehicle	12	(0.7)
Intentional neglect	1	(0.1)
Other (single method)	3	(0.2)
Poisoning and other method type§§	5	(0.3)
Other combination of methods§§	3	(0.2)
Unknown	24	(1.4)
Total	1,774	(100.0)

*Percentages might not total 100% because of rounding.

†Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§Asian/Pacific Islander.

¶American Indian/Alaska Native.

**Includes persons of any race.

††Includes only those decedents aged >18 years.

§§Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 33. Number* and percentage of suicides by former or current military personnel, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC) [§]	1,139	(64.2)	360	(31.6)
BAC ≤0.08 g/dL [§]			121	(33.6)
BAC >0.08 g/dL [§]			231	(64.2)
Alcohol positive, level unknown			8	(2.2)
Amphetamines	638	(36.0)	15	(2.4)
Antidepressants	561	(31.6)	136	(24.2)
Cocaine	672	(37.9)	35	(5.2)
Marijuana	511	(28.8)	21	(4.1)
Opiates	669	(37.7)	135	(20.2)
Other drug(s)	593	(33.4)	292	(49.2)

* N = 1,774.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 34. Number* and percentage† of suicides among former or current military personnel, by associated circumstances — National Violent Death Reporting System, 16 states,§ 2007

Circumstance	No.	(%)
Mental health/Substance abuse		
Current depressed mood	659	(43.2)
Current mental health problem	561	(36.8)
Current mental health treatment	418	(27.4)
Alcohol problem	250	(16.4)
Other substance abuse problem	121	(7.9)
Interpersonal		
Intimate partner problem	384	(25.2)
Other relationship problem (nonintimate)	127	(8.3)
Suicide of family member or friend within past 5 years	29	(1.9)
Other death of family member or friend within past 5 years	109	(7.1)
Perpetrator of interpersonal violence within past month	68	(4.5)
Victim of interpersonal violence within past month	3	(0.2)
Life stressor		
Crisis in past two weeks	494	(32.4)
Physical health problem	567	(37.2)
Job problem	168	(11.0)
Recent criminal legal problem	115	(7.5)
Financial problem	153	(10.0)
Non-criminal legal problem	49	(3.2)
School problem	1	(0.1)
Suicide event		
Left a suicide note	529	(34.7)
Disclosed intent to commit suicide	411	(26.9)
History of suicide attempt(s)	210	(13.8)

* N = 1,526. Circumstances were unknown for 248 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 35. Number* and percentage† of deaths caused by legal intervention, by victim's race/ethnicity, marital status, and location where injury occurred — National Violent Death Reporting System, 16 states,§ 2007

Characteristic	No.	(%)
Race/ethnicity		
White, non-Hispanic	66	(45.5)
Black, non-Hispanic	51	(35.2)
A/PI¶	1	(0.7)
AI/AN**	1	(0.7)
Hispanic††	26	(17.9)
Total	145	(100.0)
Marital status§§		
Married	30	(21.9)
Never married	63	(46.0)
Widowed	3	(2.2)
Divorced	31	(22.6)
Married, but separated	1	(0.7)
Single, not otherwise specified	6	(4.4)
Unknown	3	(2.2)
Total	137	(100.0)
Location of injury		
House	55	(37.9)
Street/Highway	46	(31.7)
Motor vehicle	5	(3.4)
Commercial/Retail area	5	(3.4)
Industrial or construction area	1	(0.7)
Parking lot/Public garage	13	(9.0)
Park, playground, sports/athletic area	4	(2.8)
Hospital or medical facility	1	(0.7)
Jail/Prison	2	(1.4)
Natural area	6	(4.1)
Hotel/Motel	1	(0.7)
Other	3	(2.1)
Unknown	3	(2.1)
Total	145	(100.0)

* No. incidents = 141; no. victim decedents = 141; no. suspect decedents = 4. Number of incidents is four more than the number provided in Table 1. Four of the "other combinations of death" included at least one legal intervention death.

† Percentages might not total 100% because of rounding.

§ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

¶ Asian/Pacific Islander.

** American Indian/Alaska Native.

†† Includes persons of any race.

§§ Includes only those victims aged >18 years.

TABLE 36. Number and percentage* of deaths caused by legal intervention, by sex and age group — National Violent Death Reporting System, 16 states,† 2007

Characteristic	Male		Female		Total	
	No.	(%)	No.	(%)	No.	(%)
Age						
<1	—	—	—	—	—	—
1–4	—	—	—	—	—	—
5–9	—	—	—	—	—	—
10–14	—	—	—	—	—	—
15–19	11	(7.7)	—	—	11	(7.6)
20–24	29	(20.4)	—	—	29	(20.0)
25–29	23	(16.2)	—	—	23	(15.9)
30–34	22	(15.5)	—	—	22	(15.2)
35–44	30	(21.10)	1	(33.3)	31	(21.4)
45–54	17	(12.0)	2	(66.7)	19	(13.10)
55–64	4	(2.8)	—	—	4	(2.8)
65–74	5	(3.5)	—	—	5	(3.4)
75–84	1	(0.7)	—	—	1	(0.7)
≥85	—	—	—	—	—	—
Total	142	(100.0)	3	(100.0)	145	(100.0)

* Percentages might not total 100% because of rounding.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

TABLE 37. Number* and percentage of deaths caused by legal intervention, by toxicology variable — National Violent Death Reporting System, 16 states,† 2007

Toxicology variable	Tested		Positive	
	No.	(%)	No.	(%)
Blood alcohol concentration (BAC)§	127	(87.6)	49	(38.6)
BAC ≤0.08 g/dL§			15	(30.6)
BAC >0.08 g/dL§			33	(67.4)
Alcohol positive, level unknown			1	(2.0)
Amphetamines	98	(67.6)	20	(20.4)
Antidepressants	74	(51.0)	4	(5.4)
Cocaine	110	(75.9)	27	(24.6)
Marijuana	72	(49.7)	22	(30.6)
Opiates	108	(74.5)	10	(9.3)
Other drug(s)	87	(60.0)	25	(28.7)

* N = 145.

† Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

§ The alcohol variable reflects the blood alcohol content (BAC) of victims using 0.08% as the standard for intoxication. The other substances are indicated if there are any positive results. The levels for these substances are not measured.

TABLE 38. Number,* percentage†, and rate‡ of suicides among persons aged ≥50 years, by age group, sex, race/ethnicity, marital status, location in which injury occurred, and method — National Violent Death Reporting System, 16 states¶, 2007

Age in Years	50–59			60–69			70–79			≥80			Total		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Sex															
Male	1,307	(74.3)	25.5	688	(76.3)	21.6	501	(85.9)	28.4	379	(87.7)	40.0	2,875	(78.2)	26.1
Female	453	(25.7)	8.4	214	(23.7)	6.0	82	(14.1)	3.6	53	(12.3)	3.0	802	(21.8)	5.1
Total	1,760	(100.0)	16.7	902	(100.0)	13.4	583	(100.0)	14.4	432	(100.0)	15.8	3,677	(100.0)	15.3
Race/ethnicity															
White, non-Hispanic	1,599	(90.9)	19.7	843	(93.5)	15.5	544	(93.3)	16.2	417	(96.5)	17.7	3,403	(92.5)	17.7
Black, non-Hispanic	65	(3.7)	4.6	25	(2.8)	3.3	14	(2.4)	NR**	7	(1.6)	NR**	111	(3.0)	3.9
A/PI††	18	(1.0)	NR**	9	(1.0)	NR**	4	(0.7)	NR**	4	(0.9)	NR**	35	(1.0)	5.6
AI/AN§§	16	(0.9)	NR**	4	(0.4)	NR**	7	(1.2)	NR**	—	—	NR**	27	(0.7)	11.7
Hispanic¶¶	58	(3.3)	10.1	20	(2.2)	6.9	13	(2.2)	NR**	4	(0.9)	NR**	95	(2.6)	8.6
Other	3	(0.2)	NR**	1	(0.1)	NR**	1	(0.2)	NR**	—	—	NR**	5	(0.1)	NR**
Unknown	1	—	NR**	—	—	NR**	—	—	NR**	—	—	NR**	1	—	NR**
Total	1,760	(100.0)	16.7	902	(100.0)	13.4	583	(100.0)	14.4	432	(100.0)	15.8	3,677	(100.0)	15.3
Marital Status															
Married	791	(44.9)	NR***	449	(49.8)	NR***	288	(49.4)	NR***	170	(39.4)	NR***	1,698	(46.2)	NR***
Never married	221	(12.6)	NR***	65	(7.2)	NR***	29	(5.0)	NR***	22	(5.1)	NR***	337	(9.2)	NR***
Widowed	87	(4.9)	NR***	88	(9.8)	NR***	150	(25.7)	NR***	200	(46.3)	NR***	525	(14.3)	NR***
Divorced	611	(34.7)	NR***	286	(31.7)	NR***	112	(19.2)	NR***	35	(8.1)	NR***	1,044	(28.4)	NR***
Married, but separated	10	(0.6)	NR***	3	(0.3)	NR***	2	(0.3)	NR***	—	—	NR***	15	(0.4)	NR***
Single, not otherwise specified	17	(1.0)	NR***	2	(0.2)	NR***	1	(0.2)	NR***	2	(0.5)	NR***	22	(0.6)	NR***
Unknown	23	(1.3)	NR***	9	(1.0)	NR***	1	(0.2)	NR***	3	(0.7)	NR***	36	(1.0)	NR***
Total	1,760	(100.0)	NR***	902	(100.0)	NR***	583	(100.0)	NR***	432	(100.0)	NR***	3,677	(100.0)	NR***
Location															
House	1,372	(78.0)	13.0	736	(81.6)	11.0	512	(87.8)	12.6	392	(90.5)	14.3	3,012	(81.9)	12.5
Street/Highway	48	(2.7)	0.5	18	(2.0)	NR**	13	(2.2)	NR**	5	(1.2)	NR**	84	(2.3)	0.3
Motor vehicle	45	(2.6)	0.4	14	(1.6)	NR**	2	(0.3)	NR**	5	(1.2)	NR**	66	(1.8)	0.3
Bar/Nightclub	1	(0.1)	NR**	—	—	NR**	—	—	NR**	—	—	NR**	1	—	NR**
Commercial/Retail area	4	(0.2)	NR**	5	(0.6)	NR**	2	(0.3)	NR**	1	(0.2)	NR**	12	(0.3)	NR**
Industrial or construction area	7	(0.4)	NR**	—	—	NR**	1	(0.2)	NR**	—	—	NR**	8	(0.2)	NR**
Office building	7	(0.4)	NR**	3	(0.3)	NR**	—	—	NR**	1	(0.2)	NR**	11	(0.3)	NR**
Parking lot/Public garage	27	(1.5)	0.3	10	(1.1)	NR**	4	(0.7)	NR**	3	(0.7)	NR**	44	(1.2)	0.2
Abandoned house, building, or warehouse	—	—	NR**	—	—	NR**	—	—	NR**	—	—	NR**	—	—	NR**
Park, playground, sports/Athletic area	21	(1.2)	0.2	14	(1.6)	NR**	8	(1.4)	NR**	2	(0.5)	NR**	45	(1.2)	0.2
Preschool/School/College/School bus	1	(0.1)	NR**	—	—	NR**	—	—	NR**	—	—	NR**	1	—	NR**
Public transportation/Station/Railroad tracks	6	(0.3)	NR**	1	(0.1)	NR**	1	(0.2)	NR**	1	(0.2)	NR**	9	(0.2)	NR**
Hospital or medical facility	9	(0.5)	NR**	5	(0.6)	NR**	2	(0.3)	NR**	2	(0.5)	NR**	18	(0.5)	NR**
Supervised residential facility	10	(0.6)	NR**	3	(0.3)	NR**	1	(0.2)	NR**	2	(0.5)	NR**	16	(0.4)	NR**
Jail/Prison	10	(0.6)	NR**	4	(0.4)	NR**	—	—	NR**	—	—	NR**	14	(0.4)	NR**
Farm	5	(0.3)	NR**	2	(0.2)	NR**	6	(1.0)	NR**	1	(0.2)	NR**	14	(0.4)	NR**
Natural area	83	(4.7)	0.8	32	(3.5)	0.5	11	(1.9)	NR**	5	(1.2)	NR**	131	(3.6)	0.5
Hotel/Motel	36	(2.0)	0.3	10	(1.1)	NR**	4	(0.7)	NR**	2	(0.5)	NR**	52	(1.4)	0.2
Other	47	(2.7)	0.4	20	(2.2)	0.3	10	(1.7)	NR**	7	(1.6)	NR**	84	(2.3)	0.3
Unknown	21	(1.2)	0.2	25	(2.8)	0.4	6	(1.0)	NR**	3	(0.7)	NR**	55	(1.5)	0.2
Total	1,760	(100.0)	16.7	902	(100.0)	13.4	583	(100.0)	14.4	432	(100.0)	15.8	3,677	(100.0)	15.3

TABLE 38. (Continued) Number,* percentaget, and rates§ of suicides among persons aged ≥50 years, by age group, sex, race/ethnicity, marital status, location in which injury occurred, and method — National Violent Death Reporting System, 16 states¶, 2007

Age in Years	50–59			60–69			70–79			≥80			Total		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	%	Rate
Method															
Firearm	877	(49.8)	8.3	569	(63.1)	8.5	449	(77.0)	11.1	332	(76.9)	12.1	2,227	(60.6)	9.3
Sharp instrument	38	(2.2)	0.4	26	(2.9)	0.4	5	(0.9)	NR**	7	(1.6)	NR**	76	(2.1)	0.3
Blunt instrument	2	(0.1)	NR**	—	—	NR**	—	—	NR**	—	—	NR**	2	(0.1)	NR**
Poisoning	470	(26.7)	4.5	164	(18.2)	2.4	57	(9.8)	1.4	31	(7.2)	1.1	722	(19.6)	3.0
Hanging/Strangulation/Suffocation	271	(15.4)	2.6	99	(11.0)	1.5	49	(8.4)	1.2	43	(10.0)	1.6	462	(12.6)	1.9
Fall	27	(1.5)	0.3	11	(1.2)	NR**	5	(0.9)	NR**	7	(1.6)	NR**	50	(1.4)	0.2
Drowning	19	(1.1)	NR**	11	(1.2)	NR**	6	(1.0)	NR**	4	(0.9)	NR**	40	(1.1)	0.2
Fire/Burns	6	(0.3)	NR**	1	(0.1)	NR**	1	(0.2)	NR**	1	(0.2)	NR**	9	(0.2)	NR**
Motor vehicle	19	(1.1)	NR**	6	(0.7)	NR**	1	(0.2)	NR**	1	(0.2)	NR**	27	(0.7)	0.1
Other (single method)	7	(0.4)	NR**	3	(0.3)	NR**	1	(0.2)	NR**	1	(0.2)	NR**	12	(0.3)	NR**
Firearm and poisoning†††	—	—	NR**	—	—	NR**	—	—	NR**	—	—	NR**	—	—	NR**
Firearm and other method type†††	—	—	NR**	1	(0.1)	NR**	—	—	NR**	—	—	NR**	1	—	NR**
Poisoning and other method type†††	5	(0.3)	NR**	3	(0.3)	NR**	1	(0.2)	NR**	1	(0.2)	NR**	10	(0.3)	NR**
Other combination of methods††	1	(0.1)	NR**	1	(0.1)	NR**	—	—	NR**	1	(0.2)	NR**	3	(0.1)	NR**
Unknown	18	(1.0)	NR**	7	(0.8)	NR**	8	(1.4)	NR**	3	(0.7)	NR**	36	(1.0)	0.1
Total	1,760	(100.0)	16.7	902	(100.0)	13.4	583	(100.0)	14.4	432	(100.0)	15.8	3,677	(100.0)	15.3

* No. incidents = 3,671; no. decedents = 3,677.

† Percentages might not total 100% because of rounding.

§ Per 100,000 population.

¶ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

** Rates not reported when number of decedents is <20.

†† Asian/Pacific Islander.

§§ American Indian/Alaskan Native.

¶¶ Includes persons of any race.

*** Rates for marital status cannot be computed because denominators are unknown.

††† Deaths involving more than one method and for which injury evidence indicates one method and for which evidence did not indicate which method caused the fatal injury.

TABLE 39. Number* and percentage† of suicides among persons aged ≥50 years, by age group and associated circumstances — National Violent Death Reporting System, 16 states,‡ 2007

Associated circumstances	50–59		60–69		70–79		≥80		Total	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Mental health/Substance abuse										
Current depressed mood	680	(38.6)	351	(38.9)	227	(38.9)	178	(41.2)	1,436	(39.1)
Current mental health problem	803	(45.6)	356	(39.5)	187	(32.1)	93	(21.5)	1,439	(39.1)
Current mental health treatment	597	(33.9)	296	(32.8)	146	(25.0)	59	(13.7)	1,098	(29.9)
Alcohol problem	348	(19.8)	117	(13.0)	34	(5.8)	8	(1.9)	507	(13.8)
Other substance abuse problem	189	(10.7)	33	(3.7)	6	(1.0)	1	(0.2)	229	(6.2)
Interpersonal										
Intimate partner problem	378	(21.5)	107	(11.9)	40	(6.9)	18	(4.2)	543	(14.8)
Other relationship problem (nonintimate)	164	(9.3)	71	(7.9)	25	(4.3)	12	(2.8)	272	(7.4)
Suicide of family member or friend during previous past 5 years	23	(1.3)	9	(1.0)	6	(1.0)	6	(1.4)	44	(1.2)
Other death of family member or friend during previous 5 years	118	(6.7)	54	(6.0)	47	(8.1)	47	(10.9)	266	(7.2)
Perpetrator of interpersonal violence within previous month	63	(3.6)	25	(2.8)	12	(2.1)	6	(1.4)	106	(2.9)
Victim of interpersonal violence within previous month	3	(0.2)	—	—	—	—	—	—	3	(0.1)
Life stressor										
Crisis in past 2 weeks	405	(23.0)	172	(19.1)	127	(21.8)	78	(18.1)	782	(21.3)
Physical health problem	367	(20.9)	301	(33.4)	298	(51.1)	247	(57.2)	1,213	(33.0)
Job problem	239	(13.6)	51	(5.7)	8	(1.4)	1	(0.2)	299	(8.1)
Recent criminal legal problem	128	(7.3)	46	(5.1)	6	(1.0)	2	(0.5)	182	(4.9)
Non-criminal legal problem	59	(3.4)	13	(1.4)	3	(0.5)	4	(0.9)	79	(2.1)
Financial problem	243	(13.8)	101	(11.2)	19	(3.3)	6	(1.4)	369	(10.0)
School problem	—	—	—	—	—	—	—	—	—	—
Suicide event										
Left a suicide note	597	(33.9)	260	(28.8)	146	(25.0)	128	(29.6)	1,131	(30.8)
Disclosed intent to commit suicide	448	(25.5)	210	(23.3)	150	(25.7)	99	(22.9)	907	(24.7)
History of suicide attempt(s)	326	(18.5)	118	(13.1)	51	(8.7)	24	(5.6)	519	(14.1)

* N = 3,172. Circumstances were unknown for 505 deaths.

† Percentages might exceed 100% because multiple circumstances might have been coded.

‡ Alaska, Colorado, Georgia, Kentucky, Maryland, Massachusetts, North Carolina, New Jersey, New Mexico, Oklahoma, Oregon, Rhode Island, South Carolina, Utah, Virginia, and Wisconsin.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, visit *MMWR*'s free subscription page at <http://www.cdc.gov/mmwr/mmwrsubscribe.html>. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data presented by the Notifiable Disease Data Team and 122 Cities Mortality Data Team in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. Address all inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to mmwrq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.